

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 April 2006 (27.04.2006)

PCT

(10) International Publication Number
WO 2006/044686 A2

(51) International Patent Classification: **Not classified**

(21) International Application Number:
PCT/US2005/037058

(22) International Filing Date: 17 October 2005 (17.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/967,879 18 October 2004 (18.10.2004) US
11/141,288 31 May 2005 (31.05.2005) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF WASHINGTON** [US/US]; 4311 - 11th Avenue NE, Suite 500, Seattle, WA 98105-4608 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **RIEDER, Mark, J.** [US/US]; 1234 NE 98th Street, Seattle, WA 98115 (US).
RETTIE, Allan [GB/US]; 5777 Captain Vancouver Drive, Langley, WA 98260 (US).

(74) Agents: **ARENSEN, Tanya, A.** et al.; Medlen & Carroll, LLP, 101 Howard Street, Suite 350, San Francisco, CA 94105 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **METHODS AND COMPOSITIONS FOR PREDICTING DRUG RESPONSES**

(57) Abstract: The present invention relates to methods and compositions for predicting drug responses. In particular, the present invention provides methods and compositions for determining individualized Warfarin dosages based on the presence or absence of polymorphisms in the *VKORC1* gene. The present invention further provides methods and compositions for determining individualized Warfarin dosages based on the level of expression of the *VKORC1* gene.

WO 2006/044686 A2

METHODS AND COMPOSITIONS FOR PREDICTING DRUG RESPONSES

This application was supported in part by NHLBI – Program for Genomic Applications (PGA) grant (U01 HL66682), Program for Genomic Applications (PGA) grant
5 U01 HL66682, NIH General Medical Sciences grant GM068797 and UW NIEHS sponsored Center for Ecogenetics and Environmental Health, grant NIEHS P30ES07033. The government has certain rights in the invention.

FIELD OF THE INVENTION

10 The present invention relates to methods and compositions for predicting drug responses. In particular, the present invention provides methods and compositions for determining individualized Warfarin dosages based on the presence or absence of polymorphisms in the *VKORC1* gene. The present invention further provides methods and
15 compositions for determining individualized Warfarin dosages based on the level of expression of the *VKORC1* gene.

BACKGROUND OF THE INVENTION

More than 3 billion prescriptions are written each year in the U.S. alone, effectively preventing or treating illness in hundreds of millions of people. But prescription
20 medications also can cause powerful toxic effects in a patient. These effects are called adverse drug reactions (ADR). Adverse drug reactions can cause serious injury and or even death. Differences in the ways in which individuals utilize and eliminate drugs from their bodies are one of the most important causes of ADRs. Differences in metabolism also cause doses of drugs to be less effective than desired in some individuals.

25 More than 106,000 Americans die – three times as many as are killed in automobile accidents – and an additional 2.1 million are seriously injured every year due to adverse drug reactions. ADRs are the fourth leading cause of death for Americans. Only heart disease, cancer and stroke cause more deaths each year. Seven percent of all hospital patients are affected by serious or fatal ADRs. More than two-thirds of all ADRs occur
30 outside hospitals. Adverse drug reactions are a severe, common and growing cause of death, disability and resource consumption.

It is estimated that drug-related anomalies account for nearly 10 percent of all hospital admissions. Drug-related morbidity and mortality in the U.S. is estimated to cost from \$76.6 to \$136 billion annually.

Most prescription drugs are currently prescribed at standard doses in a "one size fits all" method. This "one size fits all" method, however, does not consider important genetic differences that give different individuals dramatically different abilities to metabolize and derive benefit from a particular drug. Genetic differences may be influenced by race or ethnicity, but may also be largely unpredictable without identifying correlating genomics. What is needed are improved methods for predicting an individual's response to a given drug or a particular dosage of a drug.

SUMMARY OF THE INVENTION

The present invention relates to methods and compositions for predicting drug responses. In particular, the present invention provides methods and compositions for determining individualized Warfarin dosages based on the presence or absence of polymorphisms in the *VKORC1* gene. The present invention further provides methods and compositions for determining individualized Warfarin dosages based on the level of expression of the *VKORC1* gene.

Accordingly, in some embodiments, the present invention provides compositions, kits, and methods for determining the level of the presence of *VKORC1* polymorphisms and/or gene expression (e.g., by measuring *VKORC1* mRNA or protein expression) and correlating expression levels and/or polymorphisms with Warfarin dosages.

Accordingly, in some embodiments, the present invention provides a method, comprising the steps of: providing a sample from a subject; and determining the subject's *VKORC1* haplotype, SNP genotype, or SNP in linkage disequilibrium with any diagnostic SNP. In some embodiments, the method further comprises the step of determining the subject's optimal Warfarin dose based on the subject's *VKORC1* haplotype (e.g., H1, H2, H7, H8, or H9 haplotypes). In some embodiments, the method further comprises the step of determining the subject's CYP2C9 genotype. In some embodiments, determining the subject's *VKORC1* genotype comprises the use of a nucleic acid based detection assay (e.g., a sequencing assay or a hybridization assay). In some embodiments, the method further comprises the step of determining the subject's Clade type (e.g., AA, AB, or BB Clade types).

In other embodiments, the present invention provides a method, comprising the steps of providing a sample from a subject; detecting the genotype of a single nucleotide polymorphism at one or more positions of SEQ ID NO:1 (e.g., positions 381, 3673, 5808, 6484, 6853, 7566, and 9041 or any polymorphism in linkage disequilibrium with these

sites); and determining the subject's optimal Warfarin dosage based on said genotype of the single nucleotide polymorphism. In some embodiments, determining the subject's *VKORC1* genotype comprises the use of a nucleic acid based detection assay (e.g., a sequencing assay or a hybridization assay).

5 The present invention further provides a kit for determining a subject's optimal dose of a blood clotting drug (e.g., Warfarin), comprising: a detection assay, wherein the detection assay is capable of specifically detecting the subject's *VKORC1* haplotype (e.g., H1, H2, H7, H8, or H9 haplotypes); and instructions for determining the subject's optimal Warfarin dosage. In some embodiments, the kit further comprises reagents for determining
10 the subject's CYP2C9 genotype. In some embodiments, the detection assay is a nucleic acid based detection assay (e.g., a sequencing assay or a hybridization assay). In some embodiments, the kit further comprises instructions for determining the subject's Clade type (e.g., AA, AB, or BB Clade types).

 In further embodiments, the present invention provides a method, comprising:
15 providing a sample from a subject; and determining the subject's *VKORC1* expression level to determine responsiveness to Warfarin therapy. In some embodiments, the method further comprised the step of determining the subject's optimal Warfarin dose based on the subject's *VKORC1* expression level. In certain embodiments, the method further comprises the step of determining said subject's CYP2C9 genotype. In some embodiments, determining the
20 subject's *VKORC1* expression level comprises determining the amount of *VKORC1* mRNA expressed by said subject (e.g., by using a quantitative RT-PCR assay or a nucleic acid hybridization assay). In other embodiments, determining the subject's *VKORC1* expression level comprises determining the amount of *VKORC1* polypeptide expressed by the subject (e.g., by exposing the sample to an antibody that specifically binds to the *VKORC1*
25 polypeptide).

 The present invention further provides a kit for determining a subject optimal Warfarin dosage, comprising: reagents for performing a detection assay, wherein the detection assay is configured to specifically detect the subject's *VKORC1* expression level; and instructions for determining the subject's optimal Warfarin dosage. In some
30 embodiments, the reagents comprise reagents for determining the amount of *VKORC1* mRNA expressed by the subject (e.g., reagents for a quantitative RT-PCR assay or a nucleic acid hybridization assay). In other embodiments, the reagents comprise reagents for determining the amount of *VKORC1* polypeptide expressed by the subject (e.g., an antibody

that specifically binds to the *VKORC1* polypeptide). In some embodiments, the kit further comprises reagents for determining said subject's *CYP2C9* genotype.

DESCRIPTION OF THE FIGURES

5 Figure 1 shows the effect of *VKORC1* genealogic clades on clinical warfarin dose. The upper panel shows common haplotypes determined from *VKORC1* (H1 (SEQ ID NO:14), H2 (SEQ ID NO:15), H7 (SEQ ID NO:16), H8 (SEQ ID NO:17), and H9 (SEQ ID NO:18). The lower panel shows Warfarin dosages for clinical patients (n = 185) classified according to known functional mutations at the *CYP2C9* locus and *VKORC1* Clade (A/A (white bars), A/B (grey bars), and B/B (black bars)).

10 Figure 2 shows the nucleic acid sequence of the extended genomic reference sequence for the *VKORC1* (SEQ ID NO:1) gene.

Figure 3 shows that *VKORC1* haplotype groups correlate with mRNA expression.

15 DEFINITIONS

To facilitate an understanding of the present invention, a number of terms and phrases are defined below:

As used herein, the term "single nucleotide polymorphism" or "SNP", refers to any position along a nucleotide sequence that has one or more variant nucleotides. Single
20 nucleotide polymorphisms (SNPs) are the most common form of DNA sequence variation found in the human genome and are generally defined as a difference from the baseline reference DNA sequence which has been produced as part of the Human Genome Project or as a difference found between a subset of individuals drawn from the population at large. SNPs occur at an average rate of approximately 1 SNP/1000 base pairs when comparing
25 any two randomly chosen human chromosomes. Extremely rare SNPs can be identified which may be restricted to a specific individual or family, or conversely can be found to be extremely common in the general population (present in many unrelated individuals). SNPs can arise due to errors in DNA replication (i.e., spontaneously) or due to mutagenic agents (i.e., from a specific DNA damaging material) and can be transmitted during reproduction
30 of the organism to subsequent generations of individuals.

As used herein, the term "linkage disequilibrium" refers to single nucleotide polymorphisms where the genotypes are correlated between these polymorphisms. Several statistical measures can be used to quantify this relationship (i.e. D' , r^2 , etc) reference (See e.g., Devlin and Risch 1995 Sep 20;29(2):311-22). In some embodiments, a SNP-SNP

pair is considered to be in linkage disequilibrium if $r^2 > 0.5$,

As used herein, the term "haplotype" refers to a group of closely linked alleles that are inherited together.

As used herein, the term "haplotype clade" or "clade" refers to any group of
5 haplotypes that are all more similar to one another than any of them is to any other haplotype. Clades may be identified, for example, by performing statistical cluster analysis.

As used herein, the term "subject" refers to any animal (e.g., a mammal), including, but not limited to, humans, non-human primates, rodents, and the like. Typically, the terms "subject" and "patient" are used interchangeably herein in reference to a human subject.

10 As used herein, the term "non-human transgenic animal" refers to a non-human animal (preferable a mammal, more preferably a mouse) whose endogenous *VKORC1* gene has been inactivated (e.g., as the result of a "*VKORC1*" or a "*VKORC1* knock-in") or altered (e.g., contains a polymorphic form of the *VKORC1* gene).

As used herein, the term "non-human animals" refers to all non-human animals
15 including, but are not limited to, vertebrates such as rodents, non-human primates, ovines, bovines, ruminants, lagomorphs, porcines, caprines, equines, canines, felines, aves, etc.

As used herein, the term "gene transfer system" refers to any means of delivering a composition comprising a nucleic acid sequence to a cell or tissue. For example, gene transfer systems include, but are not limited to, vectors (e.g., retroviral, adenoviral, adeno-
20 associated viral, and other nucleic acid-based delivery systems), microinjection of naked nucleic acid, polymer-based delivery systems (e.g., liposome-based and metallic particle-based systems), biolistic injection, and the like. As used herein, the term "viral gene transfer system" refers to gene transfer systems comprising viral elements (e.g., intact viruses, modified viruses and viral components such as nucleic acids or proteins) to
25 facilitate delivery of the sample to a desired cell or tissue. As used herein, the term "adenovirus gene transfer system" refers to gene transfer systems comprising intact or altered viruses belonging to the family Adenoviridae.

As used herein, the term "site-specific recombination target sequences" refers to nucleic acid sequences that provide recognition sequences for recombination factors and the
30 location where recombination takes place.

As used herein, the term "nucleic acid molecule" refers to any nucleic acid containing molecule, including but not limited to, DNA or RNA. The term encompasses sequences that include any of the known base analogs of DNA and RNA including, but not limited to, 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine,

pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethylaminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

The term "gene" refers to a nucleic acid (*e.g.*, DNA) sequence that comprises coding sequences necessary for the production of a polypeptide, precursor, or RNA (*e.g.*, rRNA, tRNA). The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (*e.g.*, enzymatic activity, ligand binding, signal transduction, immunogenicity, etc.) of the full-length or fragment are retained. The term also encompasses the coding region of a structural gene and the sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 5 kb or more on either end such that the gene corresponds to the length of the full-length mRNA. Sequences located 5' of the coding region and present on the mRNA are referred to as 5' untranslated sequences. Sequences located 3' or downstream of the coding region and present on the mRNA are referred to as 3' untranslated sequences. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are segments of a gene that are transcribed into nuclear RNA (hnRNA); introns may contain regulatory elements such as enhancers. Introns are removed or "spliced out" from the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

As used herein, the term "heterologous gene" refers to a gene that is not in its natural environment. For example, a heterologous gene includes a gene from one species introduced into another species. A heterologous gene also includes a gene native to an

organism that has been altered in some way (*e.g.*, mutated, added in multiple copies, linked to non-native regulatory sequences, etc). Heterologous genes are distinguished from endogenous genes in that the heterologous gene sequences are typically joined to DNA sequences that are not found naturally associated with the gene sequences in the chromosome or are associated with portions of the chromosome not found in nature (*e.g.*, genes expressed in loci where the gene is not normally expressed).

As used herein, the term "transgene" refers to a heterologous gene that is integrated into the genome of an organism (*e.g.*, a non-human animal) and that is transmitted to progeny of the organism during sexual reproduction.

As used herein, the term "transgenic organism" refers to an organism (*e.g.*, a non-human animal) that has a transgene integrated into its genome and that transmits the transgene to its progeny during sexual reproduction.

As used herein, the term "gene expression" refers to the process of converting genetic information encoded in a gene into RNA (*e.g.*, mRNA, rRNA, tRNA, or snRNA) through "transcription" of the gene (*i.e.*, via the enzymatic action of an RNA polymerase), and for protein encoding genes, into protein through "translation" of mRNA. Gene expression can be regulated at many stages in the process. "Up-regulation" or "activation" refers to regulation that increases the production of gene expression products (*i.e.*, RNA or protein), while "down-regulation" or "repression" refers to regulation that decreases production. Molecules (*e.g.*, transcription factors) that are involved in up-regulation or down-regulation are often called "activators" and "repressors," respectively.

In addition to containing introns, genomic forms of a gene may also include sequences located on both the 5' and 3' end of the sequences that are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (these flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5' flanking region may contain regulatory sequences such as promoters and enhancers that control or influence the transcription of the gene. The 3' flanking region may contain sequences that direct the termination of transcription, post-transcriptional cleavage and polyadenylation.

The term "wild-type" refers to a gene or gene product isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the term "modified" or "mutant" refers to a gene or gene product that displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when

compared to the wild-type gene or gene product. It is noted that naturally occurring mutants can be isolated; these are identified by the fact that they have altered characteristics (including altered nucleic acid sequences) when compared to the wild-type gene or gene product.

5 As used herein, the terms "nucleic acid molecule encoding," "DNA sequence encoding," and "DNA encoding" refer to the order or sequence of deoxyribonucleotides along a strand of deoxyribonucleic acid. The order of these deoxyribonucleotides determines the order of amino acids along the polypeptide (protein) chain. The DNA sequence thus codes for the amino acid sequence.

10 As used herein, the terms "an oligonucleotide having a nucleotide sequence encoding a gene" and "polynucleotide having a nucleotide sequence encoding a gene," means a nucleic acid sequence comprising the coding region of a gene or in other words the nucleic acid sequence that encodes a gene product. The coding region may be present in a cDNA, genomic DNA or RNA form. When present in a DNA form, the oligonucleotide or
15 polynucleotide may be single-stranded (*i.e.*, the sense strand) or double-stranded. Suitable control elements such as enhancers/promoters, splice junctions, polyadenylation signals, etc. may be placed in close proximity to the coding region of the gene if needed to permit proper initiation of transcription and/or correct processing of the primary RNA transcript. Alternatively, the coding region utilized in the expression vectors of the present invention
20 may contain endogenous enhancers/promoters, splice junctions, intervening sequences, polyadenylation signals, etc. or a combination of both endogenous and exogenous control elements.

 As used herein, the term "oligonucleotide," refers to a short length of single-stranded polynucleotide chain. Oligonucleotides are typically less than 200 residues long (*e.g.*,
25 between 15 and 100), however, as used herein, the term is also intended to encompass longer polynucleotide chains. Oligonucleotides are often referred to by their length. For example a 24 residue oligonucleotide is referred to as a "24-mer". Oligonucleotides can form secondary and tertiary structures by self-hybridizing or by hybridizing to other polynucleotides. Such structures can include, but are not limited to, duplexes, hairpins, cruciforms, bends, and triplexes.
30

 As used herein, the terms "complementary" or "complementarity" are used in reference to polynucleotides (*i.e.*, a sequence of nucleotides) related by the base-pairing rules. For example, for the sequence "5'-A-G-T-3'," is complementary to the sequence "3'-T-C-A-5'." Complementarity may be "partial," in which only some of the nucleic acids'

bases are matched according to the base pairing rules. Or, there may be "complete" or "total" complementarity between the nucleic acids. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, as well as detection methods that depend upon binding between nucleic acids.

The term "homology" refers to a degree of complementarity. There may be partial homology or complete homology (*i.e.*, identity). A partially complementary sequence is a nucleic acid molecule that at least partially inhibits a completely complementary nucleic acid molecule from hybridizing to a target nucleic acid is "substantially homologous." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization and the like) under conditions of low stringency. A substantially homologous sequence or probe will compete for and inhibit the binding (*i.e.*, the hybridization) of a completely homologous nucleic acid molecule to a target under conditions of low stringency. This is not to say that conditions of low stringency are such that non-specific binding is permitted; low stringency conditions require that the binding of two sequences to one another be a specific (*i.e.*, selective) interaction. The absence of non-specific binding may be tested by the use of a second target that is substantially non-complementary (*e.g.*, less than about 30% identity); in the absence of non-specific binding the probe will not hybridize to the second non-complementary target.

When used in reference to a double-stranded nucleic acid sequence such as a cDNA or genomic clone, the term "substantially homologous" refers to any probe that can hybridize to either or both strands of the double-stranded nucleic acid sequence under conditions of low stringency as described above.

A gene may produce multiple RNA species that are generated by differential splicing of the primary RNA transcript. cDNAs that are splice variants of the same gene will contain regions of sequence identity or complete homology (representing the presence of the same exon or portion of the same exon on both cDNAs) and regions of complete non-identity (for example, representing the presence of exon "A" on cDNA 1 wherein cDNA 2 contains exon "B" instead). Because the two cDNAs contain regions of sequence identity they will both hybridize to a probe derived from the entire gene or portions of the gene containing sequences found on both cDNAs; the two splice variants are therefore substantially homologous to such a probe and to each other.

When used in reference to a single-stranded nucleic acid sequence, the term "substantially homologous" refers to any probe that can hybridize (*i.e.*, it is the complement of) the single-stranded nucleic acid sequence under conditions of low stringency as described above.

5 As used herein, the term "hybridization" is used in reference to the pairing of complementary nucleic acids. Hybridization and the strength of hybridization (*i.e.*, the strength of the association between the nucleic acids) is impacted by such factors as the degree of complementary between the nucleic acids, stringency of the conditions involved, the T_m of the formed hybrid, and the G:C ratio within the nucleic acids. A single molecule
10 that contains pairing of complementary nucleic acids within its structure is said to be "self-hybridized."

As used herein, the term " T_m " is used in reference to the "melting temperature." The melting temperature is the temperature at which a population of double-stranded nucleic acid molecules becomes half dissociated into single strands. The equation for calculating
15 the T_m of nucleic acids is well known in the art. As indicated by standard references, a simple estimate of the T_m value may be calculated by the equation: $T_m = 81.5 + 0.41(\% G + C)$, when a nucleic acid is in aqueous solution at 1 M NaCl (See *e.g.*, Anderson and Young, Quantitative Filter Hybridization, in Nucleic Acid Hybridization [1985]). Other references include more sophisticated computations that take structural as well as sequence
20 characteristics into account for the calculation of T_m .

As used herein the term "stringency" is used in reference to the conditions of temperature, ionic strength, and the presence of other compounds such as organic solvents, under which nucleic acid hybridizations are conducted. Under "low stringency conditions" a nucleic acid sequence of interest will hybridize to its exact complement, sequences with
25 single base mismatches, closely related sequences (*e.g.*, sequences with 90% or greater homology), and sequences having only partial homology (*e.g.*, sequences with 50-90% homology). Under "medium stringency conditions," a nucleic acid sequence of interest will hybridize only to its exact complement, sequences with single base mismatches, and closely related sequences (*e.g.*, 90% or greater homology). Under "high stringency conditions," a
30 nucleic acid sequence of interest will hybridize only to its exact complement, and (depending on conditions such as temperature) sequences with single base mismatches. In other words, under conditions of high stringency the temperature can be raised so as to exclude hybridization to sequences with single base mismatches.

"High stringency conditions" when used in reference to nucleic acid hybridization comprise conditions equivalent to binding or hybridization at 42°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄·H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.5% SDS, 5X Denhardt's reagent and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 0.1X SSPE, 1.0% SDS at 42°C when a probe of about 500 nucleotides in length is employed.

"Medium stringency conditions" when used in reference to nucleic acid hybridization comprise conditions equivalent to binding or hybridization at 42°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄·H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.5% SDS, 5X Denhardt's reagent and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 1.0X SSPE, 1.0% SDS at 42°C when a probe of about 500 nucleotides in length is employed.

"Low stringency conditions" comprise conditions equivalent to binding or hybridization at 42°C in a solution consisting of 5X SSPE (43.8 g/l NaCl, 6.9 g/l NaH₂PO₄·H₂O and 1.85 g/l EDTA, pH adjusted to 7.4 with NaOH), 0.1% SDS, 5X Denhardt's reagent [50X Denhardt's contains per 500 ml: 5 g Ficoll (Type 400, Pharmacia), 5 g BSA (Fraction V; Sigma)] and 100 µg/ml denatured salmon sperm DNA followed by washing in a solution comprising 5X SSPE, 0.1% SDS at 42°C when a probe of about 500 nucleotides in length is employed.

The art knows well that numerous equivalent conditions may be employed to comprise low stringency conditions; factors such as the length and nature (DNA, RNA, base composition) of the probe and nature of the target (DNA, RNA, base composition, present in solution or immobilized, etc.) and the concentration of the salts and other components (*e.g.*, the presence or absence of formamide, dextran sulfate, polyethylene glycol) are considered and the hybridization solution may be varied to generate conditions of low stringency hybridization different from, but equivalent to, the above listed conditions. In addition, the art knows conditions that promote hybridization under conditions of high stringency (*e.g.*, increasing the temperature of the hybridization and/or wash steps, the use of formamide in the hybridization solution, etc.) (see definition above for "stringency").

As used herein, the term "detection assay" refers to an assay for detecting the presence or absence of variant nucleic acid sequences (*e.g.*, polymorphism or mutations) in a given allele of a particular gene (*e.g.*, the *VKORC1* gene).

The term "isolated" when used in relation to a nucleic acid, as in "an isolated oligonucleotide" or "isolated polynucleotide" refers to a nucleic acid sequence that is identified and separated from at least one component or contaminant with which it is ordinarily associated in its natural source. Isolated nucleic acid is such present in a form or setting that is different from that in which it is found in nature. In contrast, non-isolated nucleic acids as nucleic acids such as DNA and RNA found in the state they exist in nature. For example, a given DNA sequence (*e.g.*, a gene) is found on the host cell chromosome in proximity to neighboring genes; RNA sequences, such as a specific mRNA sequence encoding a specific protein, are found in the cell as a mixture with numerous other mRNAs that encode a multitude of proteins. However, isolated nucleic acid encoding a given protein includes, by way of example, such nucleic acid in cells ordinarily expressing the given protein where the nucleic acid is in a chromosomal location different from that of natural cells, or is otherwise flanked by a different nucleic acid sequence than that found in nature. The isolated nucleic acid, oligonucleotide, or polynucleotide may be present in single-stranded or double-stranded form. When an isolated nucleic acid, oligonucleotide or polynucleotide is to be utilized to express a protein, the oligonucleotide or polynucleotide will contain at a minimum the sense or coding strand (*i.e.*, the oligonucleotide or polynucleotide may be single-stranded), but may contain both the sense and anti-sense strands (*i.e.*, the oligonucleotide or polynucleotide may be double-stranded).

As used herein, the term "purified" or "to purify" refers to the removal of components (*e.g.*, contaminants) from a sample. For example, antibodies are purified by removal of contaminating non-immunoglobulin proteins; they are also purified by the removal of immunoglobulin that does not bind to the target molecule. The removal of non-immunoglobulin proteins and/or the removal of immunoglobulins that do not bind to the target molecule results in an increase in the percent of target-reactive immunoglobulins in the sample. In another example, recombinant polypeptides are expressed in bacterial host cells and the polypeptides are purified by the removal of host cell proteins; the percent of recombinant polypeptides is thereby increased in the sample.

"Amino acid sequence" and terms such as "polypeptide" or "protein" are not meant to limit the amino acid sequence to the complete, native amino acid sequence associated with the recited protein molecule.

The term "native protein" as used herein to indicate that a protein does not contain amino acid residues encoded by vector sequences; that is, the native protein contains only

those amino acids found in the protein as it occurs in nature. A native protein may be produced by recombinant means or may be isolated from a naturally occurring source.

As used herein the term "portion" when in reference to a protein (as in "a portion of a given protein") refers to fragments of that protein. The fragments may range in size from
5 four amino acid residues to the entire amino acid sequence minus one amino acid.

As used herein, the term "vector" is used in reference to nucleic acid molecules that transfer DNA segment(s) from one cell to another. The term "vehicle" is sometimes used interchangeably with "vector." Vectors are often derived from plasmids, bacteriophages, or plant or animal viruses.

10 The term "expression vector" as used herein refers to a recombinant DNA molecule containing a desired coding sequence and appropriate nucleic acid sequences necessary for the expression of the operably linked coding sequence in a particular host organism. Nucleic acid sequences necessary for expression in prokaryotes usually include a promoter, an operator (optional), and a ribosome binding site, often along with other sequences.
15 Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

The terms "overexpression" and "overexpressing" and grammatical equivalents, are used in reference to levels of mRNA to indicate a level of expression approximately 3-fold higher (or greater) than that observed in a given tissue in a control or non-transgenic animal.
20 Levels of mRNA are measured using any of a number of techniques known to those skilled in the art including, but not limited to Northern blot analysis. Appropriate controls are included on the Northern blot to control for differences in the amount of RNA loaded from each tissue analyzed (*e.g.*, the amount of 28S rRNA, an abundant RNA transcript present at essentially the same amount in all tissues, present in each sample can be used as a means of
25 normalizing or standardizing the mRNA-specific signal observed on Northern blots). The amount of mRNA present in the band corresponding in size to the correctly spliced transgene RNA is quantified; other minor species of RNA which hybridize to the transgene probe are not considered in the quantification of the expression of the transgenic mRNA.

The term "transfection" as used herein refers to the introduction of foreign DNA into
30 eukaryotic cells. Transfection may be accomplished by a variety of means known to the art including calcium phosphate-DNA co-precipitation, DEAE-dextran-mediated transfection, polybrene-mediated transfection, electroporation, microinjection, liposome fusion, lipofection, protoplast fusion, retroviral infection, and biolistics.

The term "stable transfection" or "stably transfected" refers to the introduction and integration of foreign DNA into the genome of the transfected cell. The term "stable transfectant" refers to a cell that has stably integrated foreign DNA into the genomic DNA.

5 The term "transient transfection" or "transiently transfected" refers to the introduction of foreign DNA into a cell where the foreign DNA fails to integrate into the genome of the transfected cell. The foreign DNA persists in the nucleus of the transfected cell for several days. During this time the foreign DNA is subject to the regulatory controls that govern the expression of endogenous genes in the chromosomes. The term "transient transfectant" refers to cells that have taken up foreign DNA but have failed to integrate this
10 DNA.

As used, the term "eukaryote" refers to organisms distinguishable from "prokaryotes." It is intended that the term encompass all organisms with cells that exhibit the usual characteristics of eukaryotes, such as the presence of a true nucleus bounded by a nuclear membrane, within which lie the chromosomes, the presence of membrane-bound
15 organelles, and other characteristics commonly observed in eukaryotic organisms. Thus, the term includes, but is not limited to such organisms as fungi, protozoa, and animals (e.g., humans).

As used herein, the term "*in vitro*" refers to an artificial environment and to processes or reactions that occur within an artificial environment. In vitro environments can
20 consist of, but are not limited to, test tubes and cell culture. The term "*in vivo*" refers to the natural environment (e.g., an animal or a cell) and to processes or reaction that occur within a natural environment.

The terms "test compound" and "candidate compound" refer to any chemical entity, pharmaceutical, drug, and the like that is a candidate for use to treat or prevent a disease,
25 illness, sickness, or disorder of bodily function (e.g., cancer). Test compounds comprise both known and potential therapeutic compounds. A test compound can be determined to be therapeutic by screening using the screening methods of the present invention.

As used herein, the term "sample" is used in its broadest sense. In one sense, it is meant to include a specimen or culture obtained from any source, as well as biological and
30 environmental samples. Biological samples may be obtained from animals (including humans) and encompass fluids, solids, tissues, and gases. Biological samples include blood products, such as plasma, serum and the like. Environmental samples include environmental material such as surface matter, soil, water and industrial samples. Such

examples are not however to be construed as limiting the sample types applicable to the present invention.

DETAILED DESCRIPTION OF THE INVENTION

5 Coumarin anticoagulant drugs are the definitive treatment world-wide for the long-term prevention of thromboembolic events. In 2003, a total of 21.2 million prescriptions were written for the oral anticoagulant warfarin in the United States alone. Unfortunately, warfarin poses considerable dose management problems due to a multitude of factors that can modify the anticoagulant effect of the drug: its narrow therapeutic range, discrete ethnic
10 differences in dose requirements and wide inter-individual variability in dosing. These challenges may contribute to the general under-utilization of anticoagulant therapy, particularly in stroke prevention (Fang et al., Arch Intern Med 164, 55-60 (2004); Gage et al., Stroke 31, 822-7 (2000)). Structural gene mutations in cytochrome P450 (*CYP*) 2C9, the major catabolic enzyme for the more active (*S*)-enantiomer of warfarin, are a risk factor
15 for adverse outcomes during therapy (Higashi et al., Jama 287, 1690-8 (2002)), and extremely rare mutations in *VKORC1* underlie overt warfarin resistance (Rost et al., Nature 427, 537-41 (2004)). The association of a single *VKORC1* polymorphism with Warfarin dosage has been described (D'Andrea, Blood, September 9, 2004). However, prior to the present invention, much of the variance in warfarin dose requirement remained unexplained
20 (Gage et al., Thromb Haemost 91, 87-94 (2004)).

 Warfarin exerts its antithrombotic effects by inhibiting regeneration of an essential component of clotting factor synthesis - vitamin K_{H2} (reduced vitamin K)- from vitamin K epoxide (Suttie, Adv Exp Med Biol 214, 3-16 (1987)). This enzyme activity is determined by the recently discovered vitamin K epoxide reductase gene, *VKORC1* (Li et al., Nature
25 427, 541-4 (2004); Rost et al., supra).

 Experiments conducted during the course of development of the present invention demonstrated a correlation between certain *VKORC1* haplotypes and optimal warfarin dosage. Accordingly, in some embodiments, the present invention provides methods and compositions for determining a subject's optimal Warfarin dose, as well as for related drugs
30 (e.g., drugs that involve the same biological pathway).

 Further experiments conducted during the course of development of the present invention demonstrated a correlation between *VKORC1* expression and *VKORC1* haplotypes (See e.g., Example 2). The associations between (i) the A haplotype and reduced mRNA expression and (ii) the B haplotype and increased mRNA expression

parallels the effect of these haplotypes on warfarin dose, as would be predicted by a simple, non-competitive model of enzyme inhibition by this anticoagulant (Fasco et al., Biochemistry 1983; 22:5655-60). The present invention is not limited to a particular mechanism. Indeed, an understanding of the mechanism is not necessary to practice the present invention. Nonetheless, it is contemplated that the level of *VKORC1* mRNA is directed by each haplotype and determines the level of protein synthesis of the vitamin K epoxide reductase complex which, in turn, accounts for differences in warfarin maintenance dose in patients. The primary SNP candidates that explain this effect are those that designate the major haplotype split (sites 381, 3673, 6484, 6853, and 7566) and predict warfarin maintenance dose. These SNPs were mapped to homologous regions in rat, mouse, and dog species, to identify potentially conserved non-coding sequences that encompass these sites. Only two SNPs (6484 and 6583) from the informative group are conserved; these flank exon 2 but fall outside the canonical regions required for exon splicing. It is contemplated that these regions act as regulatory sequences that bind transcription factor binding sites.

The merits of genotyping prior to, or concomitant with, treatment involving drugs like warfarin, irinotecan and thiopurine—the effectiveness of which depend on genetic variants of CYP2C9 (and now *VKORC1*), UGT1A1, and TPMT—is an area of active debate between regulatory authorities and the clinical community (Lesko et al., Nat Rev Drug Discov 2004; 3:763-9). Recently published guidelines suggest initial warfarin dosing at 5 – 10 mg/day 19. However, experiments conducted during the course of development of the present invention suggest this strategy may expose low dose *VKORC1* A/A patients to unnecessarily high doses of drug. Accordingly, in some embodiments, the present invention provides methods comprising analysis of *VKORC1* expression and polymorphisms to aid in the determination of optimal Warfarin dosages.

I. Personalized Warfarin dosing

In some embodiments, the present invention provides methods of personalized Warfarin dosing comprising identifying a subject's *VKORC1* haplotype or Clade type. As described below (See Experimental Section), experiments conducted during the course of development of the present invention identified a series of *VKORC1* polymorphisms associated with optimal Warfarin dosages. Polymorphisms at seven sites (381, 3673, 5808, 6484, 6853, 7566, and 9041) of *VKORC1* were identified. The polymorphisms were found to be associated with two low-dose (2.9 and 3.0 mg/d) haplotypes (H1 and H2) and two

high-dose (6.0 and 5.5 mg/d) haplotypes (H7 and H9). Thus, the present invention provides compositions, methods, and kits for detecting such polymorphisms and haplotypes, directly or indirectly, by any method, for predicting response to Warfarin and related drugs, selecting drugs dosage, and conducting studies on drug metabolism. These polymorphisms
5 may be detected along with other polymorphisms (e.g., CYP2C9) to enhance the information available to researchers and medical practitioners.

In some embodiments, the methods of the present invention comprise identifying a subject's haplotype and determining the subject's optimal dosage range. The methods of the present invention allow for safer and thus more widespread use of Warfarin and related
10 drugs. Exemplary methods for determining *VKORC1* polymorphisms are described below.

1. Direct sequencing Assays

In some embodiments of the present invention, *VKORC1* polymorphic sequences are detected using a direct sequencing technique. In these assays, DNA samples are first
15 isolated from a subject using any suitable method. In some embodiments, the region of interest is cloned into a suitable vector and amplified by growth in a host cell (e.g., a bacteria). In other embodiments, DNA in the region of interest is amplified using PCR.

Following amplification, DNA in the region of interest (e.g., the region containing the SNP or mutation of interest) is sequenced using any suitable method, including but not
20 limited to manual sequencing using radioactive marker nucleotides, and automated sequencing. The results of the sequencing are displayed using any suitable method. The sequence is examined and the presence or absence of a given SNP or mutation is determined.

2. PCR Assay

In some embodiments of the present invention, variant sequences are detected using a PCR-based assay. In some embodiments, the PCR assay comprises the use of
oligonucleotide primers that hybridize only to the variant or wild type allele (e.g., to the region of polymorphism or mutation). Both sets of primers are used to amplify a sample of
30 DNA. If only the mutant primers result in a PCR product, then the patient has the mutant allele. If only the wild-type primers result in a PCR product, then the patient has the wild type allele.

3. Hybridization Assays

In preferred embodiments of the present invention, variant sequences are detected using a hybridization assay. In a hybridization assay, the presence or absence of a given SNP or mutation is determined based on the ability of the DNA from the sample to
5 hybridize to a complementary DNA molecule (*e.g.*, a oligonucleotide probe). A variety of hybridization assays using a variety of technologies for hybridization and detection are available. A description of a selection of assays is provided below.

a. Direct Detection of Hybridization

10 In some embodiments, hybridization of a probe to the sequence of interest (*e.g.*, a SNP or mutation) is detected directly by visualizing a bound probe (*e.g.*, a Northern or Southern assay; *See e.g.*, Ausabel *et al.* (eds.), Current Protocols in Molecular Biology, John Wiley & Sons, NY [1991]). In these assays, genomic DNA (Southern) or RNA (Northern) is isolated from a subject. The DNA or RNA is then cleaved with a series of
15 restriction enzymes that cleave infrequently in the genome and not near any of the markers being assayed. The DNA or RNA is then separated (*e.g.*, on an agarose gel) and transferred to a membrane. A labeled (*e.g.*, by incorporating a radionucleotide) probe or probes specific for the SNP or mutation being detected is allowed to contact the membrane under a condition of low, medium, or high stringency conditions. Unbound probe is removed and
20 the presence of binding is detected by visualizing the labeled probe.

b. Detection of Hybridization Using "DNA Chip" Assays

In some embodiments of the present invention, variant sequences are detected using a DNA chip hybridization assay. In this assay, a series of oligonucleotide probes are affixed
25 to a solid support. The oligonucleotide probes are designed to be unique to a given SNP or mutation. The DNA sample of interest is contacted with the DNA "chip" and hybridization is detected.

In some embodiments, the DNA chip assay is a GeneChip (Affymetrix, Santa Clara, CA; *See e.g.*, U.S. Patent Nos. 6,045,996; 5,925,525; and 5,858,659; each of which is herein
30 incorporated by reference) assay. The GeneChip technology uses miniaturized, high-density arrays of oligonucleotide probes affixed to a "chip." Probe arrays are manufactured by Affymetrix's light-directed chemical synthesis process, which combines solid-phase chemical synthesis with photolithographic fabrication techniques employed in the semiconductor industry. Using a series of photolithographic masks to define chip

exposure sites, followed by specific chemical synthesis steps, the process constructs high-density arrays of oligonucleotides, with each probe in a predefined position in the array. Multiple probe arrays are synthesized simultaneously on a large glass wafer. The wafers are then diced, and individual probe arrays are packaged in injection-molded plastic cartridges, which protect them from the environment and serve as chambers for hybridization.

The nucleic acid to be analyzed is isolated, amplified by PCR, and labeled with a fluorescent reporter group. The labeled DNA is then incubated with the array using a fluidics station. The array is then inserted into the scanner, where patterns of hybridization are detected. The hybridization data are collected as light emitted from the fluorescent reporter groups already incorporated into the target, which is bound to the probe array. Probes that perfectly match the target generally produce stronger signals than those that have mismatches. Since the sequence and position of each probe on the array are known, by complementarity, the identity of the target nucleic acid applied to the probe array can be determined.

In other embodiments, a DNA microchip containing electronically captured probes (Nanogen, San Diego, CA) is utilized (*See e.g.*, U.S. Patent Nos. 6,017,696; 6,068,818; and 6,051,380; each of which are herein incorporated by reference). Through the use of microelectronics, Nanogen's technology enables the active movement and concentration of charged molecules to and from designated test sites on its semiconductor microchip. DNA capture probes unique to a given SNP or mutation are electronically placed at, or "addressed" to, specific sites on the microchip. Since DNA has a strong negative charge, it can be electronically moved to an area of positive charge.

First, a test site or a row of test sites on the microchip is electronically activated with a positive charge. Next, a solution containing the DNA probes is introduced onto the microchip. The negatively charged probes rapidly move to the positively charged sites, where they concentrate and are chemically bound to a site on the microchip. The microchip is then washed and another solution of distinct DNA probes is added until the array of specifically bound DNA probes is complete.

A test sample is then analyzed for the presence of target DNA molecules by determining which of the DNA capture probes hybridize, with complementary DNA in the test sample (*e.g.*, a PCR amplified gene of interest). An electronic charge is also used to move and concentrate target molecules to one or more test sites on the microchip. The electronic concentration of sample DNA at each test site promotes rapid hybridization of

sample DNA with complementary capture probes (hybridization may occur in minutes). To remove any unbound or nonspecifically bound DNA from each site, the polarity or charge of the site is reversed to negative, thereby forcing any unbound or nonspecifically bound DNA back into solution away from the capture probes. A laser-based fluorescence scanner is used to detect binding,

In still further embodiments, an array technology based upon the segregation of fluids on a flat surface (chip) by differences in surface tension (ProtoGene, Palo Alto, CA) is utilized (*See e.g.*, U.S. Patent Nos. 6,001,311; 5,985,551; and 5,474,796; each of which is herein incorporated by reference). Protogene's technology is based on the fact that fluids can be segregated on a flat surface by differences in surface tension that have been imparted by chemical coatings. Once so segregated, oligonucleotide probes are synthesized directly on the chip by ink-jet printing of reagents. The array with its reaction sites defined by surface tension is mounted on a X/Y translation stage under a set of four piezoelectric nozzles, one for each of the four standard DNA bases. The translation stage moves along each of the rows of the array and the appropriate reagent is delivered to each of the reaction site. For example, the A amidite is delivered only to the sites where amidite A is to be coupled during that synthesis step and so on. Common reagents and washes are delivered by flooding the entire surface and then removing them by spinning.

DNA probes unique for the SNP or mutation of interest are affixed to the chip using Protogene's technology. The chip is then contacted with the PCR-amplified genes of interest. Following hybridization, unbound DNA is removed and hybridization is detected using any suitable method (*e.g.*, by fluorescence de-quenching of an incorporated fluorescent group).

In yet other embodiments, a "bead array" is used for the detection of polymorphisms (Illumina, San Diego, CA; *See e.g.*, PCT Publications WO 99/67641 and WO 00/39587, each of which is herein incorporated by reference). Illumina uses a BEAD ARRAY technology that combines fiber optic bundles and beads that self-assemble into an array. Each fiber optic bundle contains thousands to millions of individual fibers depending on the diameter of the bundle. The beads are coated with an oligonucleotide specific for the detection of a given SNP or mutation. Batches of beads are combined to form a pool specific to the array. To perform an assay, the BEAD ARRAY is contacted with a prepared subject sample (*e.g.*, DNA). Hybridization is detected using any suitable method.

c. Enzymatic Detection of Hybridization

In some embodiments, hybridization of a bound probe is detected using a TaqMan assay (PE Biosystems, Foster City, CA; *See e.g.*, U.S. Patent Nos. 5,962,233 and 5,538,848, each of which is herein incorporated by reference). The assay is performed during a PCR reaction. The TaqMan assay exploits the 5'-3' exonuclease activity of DNA polymerases such as AMPLITAQ DNA polymerase. A probe, specific for a given allele or mutation, is included in the PCR reaction. The probe consists of an oligonucleotide with a 5'-reporter dye (*e.g.*, a fluorescent dye) and a 3'-quencher dye. During PCR, if the probe is bound to its target, the 5'-3' nucleolytic activity of the AMPLITAQ polymerase cleaves the probe between the reporter and the quencher dye. The separation of the reporter dye from the quencher dye results in an increase of fluorescence. The signal accumulates with each cycle of PCR and can be monitored with a fluorimeter.

In still further embodiments, polymorphisms are detected using the SNP-IT primer extension assay (Orchid Biosciences, Princeton, NJ; *See e.g.*, U.S. Patent Nos. 5,952,174 and 5,919,626, each of which is herein incorporated by reference). In this assay, SNPs are identified by using a specially synthesized DNA primer and a DNA polymerase to selectively extend the DNA chain by one base at the suspected SNP location. DNA in the region of interest is amplified and denatured. Polymerase reactions are then performed using miniaturized systems called microfluidics. Detection is accomplished by adding a label to the nucleotide suspected of being at the SNP or mutation location. Incorporation of the label into the DNA can be detected by any suitable method (*e.g.*, if the nucleotide contains a biotin label, detection is via a fluorescently labelled antibody specific for biotin). Numerous other assays are known in the art.

4. Other Detection Assays

Additional detection assays that are suitable for use in the present invention include, but are not limited to, enzyme mismatch cleavage methods (*e.g.*, Variagenics, U.S. Pat. Nos. 6,110,684, 5,958,692, 5,851,770, herein incorporated by reference in their entireties); polymerase chain reaction; branched hybridization methods (*e.g.*, Chiron, U.S. Pat. Nos. 5,849,481, 5,710,264, 5,124,246, and 5,624,802, herein incorporated by reference in their entireties); rolling circle replication (*e.g.*, U.S. Pat. Nos. 6,210,884 and 6,183,960, herein incorporated by reference in their entireties); NASBA (*e.g.*, U.S. Pat. No. 5,409,818, herein incorporated by reference in its entirety); molecular beacon technology (*e.g.*, U.S. Pat. No. 6,150,097, herein incorporated by reference in its entirety); E-sensor technology (Motorola,

U.S. Pat. Nos. 6,248,229, 6,221,583, 6,013,170, and 6,063,573, herein incorporated by reference in their entireties); INVADER assay, Third Wave Technologies; *See e.g.*, U.S. Patent Nos. 5,846,717, 6,090,543; 6,001,567; 5,985,557; and 5,994,069; each of which is herein incorporated by reference; cycling probe technology (e.g., U.S. Pat. Nos. 5,403,711, 5,011,769, and 5,660,988, herein incorporated by reference in their entireties); Dade Behring signal amplification methods (e.g., U.S. Pat. Nos. 6,121,001, 6,110,677, 5,914,230, 5,882,867, and 5,792,614, herein incorporated by reference in their entireties); ligase chain reaction (Barnay Proc. Natl. Acad. Sci USA 88, 189-93 (1991)); and sandwich hybridization methods (e.g., U.S. Pat. No. 5,288,609, herein incorporated by reference in its entirety).

5. Mass Spectroscopy Assay

In some embodiments, a MassARRAY system (Sequenom, San Diego, CA.) is used to detect variant sequences (*See e.g.*, U.S. Patent Nos. 6,043,031; 5,777,324; and 5,605,798; each of which is herein incorporated by reference). DNA is isolated from blood samples using standard procedures. Next, specific DNA regions containing the mutation or SNP of interest, about 200 base pairs in length, are amplified by PCR. The amplified fragments are then attached by one strand to a solid surface and the non-immobilized strands are removed by standard denaturation and washing. The remaining immobilized single strand then serves as a template for automated enzymatic reactions that produce genotype specific diagnostic products.

Very small quantities of the enzymatic products, typically five to ten nanoliters, are then transferred to a SpectroCHIP array for subsequent automated analysis with the SpectroREADER mass spectrometer. Each spot is preloaded with light absorbing crystals that form a matrix with the dispensed diagnostic product. The MassARRAY system uses MALDI-TOF (Matrix Assisted Laser Desorption Ionization - Time of Flight) mass spectrometry. In a process known as desorption, the matrix is hit with a pulse from a laser beam. Energy from the laser beam is transferred to the matrix and it is vaporized resulting in a small amount of the diagnostic product being expelled into a flight tube. As the diagnostic product is charged when an electrical field pulse is subsequently applied to the tube they are launched down the flight tube towards a detector. The time between application of the electrical field pulse and collision of the diagnostic product with the detector is referred to as the time of flight. This is a very precise measure of the product's molecular weight, as a molecule's mass correlates directly with time of flight with smaller

molecules flying faster than larger molecules. The entire assay is completed in less than one thousandth of a second, enabling samples to be analyzed in a total of 3-5 second including repetitive data collection. The SpectroTYPER software then calculates, records, compares and reports the genotypes at the rate of three seconds per sample.

5

6. Detection of *VKORC1* Expression

In other embodiments, the level of *VKORC1* gene expression is used to determine an individual's Warfarin dose. Experiments conducted during the course of development of the present invention (See e.g., Example 2) demonstrated a correlation between *VKORC1* haplotype and level of *VKORC1* expression. Accordingly, it is contemplated that the level of *VKORC1* expression is correlated with optimal Warfarin dosage.

10

1. Detection of RNA

In some preferred embodiments, detection of *VKORC1* expression is detected by measuring the expression of corresponding mRNA in a blood sample. mRNA expression may be measured by any suitable method, including but not limited to, those disclosed below.

15

In some embodiments, RNA is detection by Northern blot analysis. Northern blot analysis involves the separation of RNA and hybridization of a complementary labeled probe.

20

In other embodiments, RNA expression is detected by enzymatic cleavage of specific structures (INVADER assay, Third Wave Technologies; See e.g., U.S. Patent Nos. 5,846,717, 6,090,543; 6,001,567; 5,985,557; and 5,994,069; each of which is herein incorporated by reference). The INVADER assay detects specific nucleic acid (e.g., RNA) sequences by using structure-specific enzymes to cleave a complex formed by the hybridization of overlapping oligonucleotide probes.

25

In still further embodiments, RNA (or corresponding cDNA) is detected by hybridization to an oligonucleotide probe). A variety of hybridization assays using a variety of technologies for hybridization and detection are available. For example, in some embodiments, TaqMan assay (PE Biosystems, Foster City, CA; See e.g., U.S. Patent Nos. 5,962,233 and 5,538,848, each of which is herein incorporated by reference) is utilized. The assay is performed during a PCR reaction. The TaqMan assay exploits the 5'-3' exonuclease activity of the AMPLITAQ GOLD DNA polymerase. A probe consisting of an oligonucleotide with a 5'-reporter dye (e.g., a fluorescent dye) and a 3'-quencher dye is

30

included in the PCR reaction. During PCR, if the probe is bound to its target, the 5'-3' nucleolytic activity of the AMPLITAQ GOLD polymerase cleaves the probe between the reporter and the quencher dye. The separation of the reporter dye from the quencher dye results in an increase of fluorescence. The signal accumulates with each cycle of PCR and can be monitored with a fluorimeter.

In yet other embodiments, reverse-transcriptase PCR (RT-PCR) is used to detect the expression of RNA. In RT-PCR, RNA is enzymatically converted to complementary DNA or "cDNA" using a reverse transcriptase enzyme. The cDNA is then used as a template for a PCR reaction. PCR products can be detected by any suitable method, including but not limited to, gel electrophoresis and staining with a DNA specific stain or hybridization to a labeled probe. In some embodiments, the quantitative reverse transcriptase PCR with standardized mixtures of competitive templates method described in U.S. Patents 5,639,606, 5,643,765, and 5,876,978 (each of which is herein incorporated by reference) is utilized.

2. Detection of Protein

In other embodiments, gene expression of *VKORC1* is detected by measuring the expression of the corresponding protein or polypeptide. Protein expression may be detected by any suitable method. In some embodiments, proteins are detected by immunohistochemistry methods known in the art. In other embodiments, proteins are detected by their binding to an antibody raised against the protein. The generation of antibodies is described below.

Antibody binding is detected by techniques known in the art (*e.g.*, radioimmunoassay, ELISA (enzyme-linked immunosorbant assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, *in situ* immunoassays (*e.g.*, using colloidal gold, enzyme or radioisotope labels, for example), Western blots, precipitation reactions, agglutination assays (*e.g.*, gel agglutination assays, hemagglutination assays, etc.), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc.

In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many methods are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

In some embodiments, an automated detection assay is utilized. Methods for the automation of immunoassays include those described in U.S. Patents 5,885,530, 4,981,785, 6,159,750, and 5,358,691, each of which is herein incorporated by reference. In some embodiments, the analysis and presentation of results is also automated. For example, in
5 some embodiments, software that generates a prognosis based on the presence or absence of a series of proteins corresponding to cancer markers is utilized.

In other embodiments, the immunoassay described in U.S. Patents 5,599,677 and 5,672,480; each of which is herein incorporated by reference.

10 II. Kits

In some embodiments, the present invention provides kits for the detection of *VKORC1* polymorphisms. In some embodiments, the kits contain reagents specific for the detection of mRNA or cDNA (*e.g.*, oligonucleotide probes or primers). In preferred
15 embodiments, the kits contain all of the components necessary to perform a detection assay, including all controls, directions for performing assays, and any necessary software for analysis and presentation of results. In some embodiments, individual probes and reagents for detection of *VKORC1* polymorphisms are provided as analyte specific reagents. In other embodiments, the kits are provided as *in vitro* diagnostics.

In other embodiments, the present invention provides kits for determining the level
20 of *VKORC1* mRNA or protein expression in a subject. For example, in some embodiments, the kits comprise reagents for performing mRNA or protein detection assays (*e.g.*, those described above).

III. Drug Screening

25 In some embodiments, the present invention provides drug screening assays (*e.g.*, to screen for anticoagulant drugs). In some embodiments, the screening methods of the present invention utilize polymorphic forms of *VKORC1*. For example, in some embodiments, the present invention provides methods of screening for compounds that alter (*e.g.*, decrease) the activity or level of expression of one or more polymorphic forms of
30 *VKORC1*. In other embodiments, the drug screening methods described below are used to screen compounds known to alter blood clotting with different polymorphic forms of *VKORC1*.

In one screening method, candidate compounds are evaluated for their ability to alter (*e.g.*, increase or decrease) *VKORC1* expression by contacting a compound with a cell

expressing *VKORC1* and then assaying for the effect of the candidate compounds on expression. In some embodiments, the effect of candidate compounds on expression of *VKORC1* is assayed for by detecting the level of *VKORC1* mRNA expressed by the cell. mRNA expression can be detected by any suitable method, including but not limited to, those disclosed herein.

In other embodiments, the effect of candidate compounds is assayed by measuring the level of *VKORC1* polypeptide. The level of polypeptide expressed can be measured using any suitable method, including but not limited to, those disclosed herein or by monitoring a phenotype (e.g., clotting speed).

In some embodiments, *in vitro* drug screens are performed using purified wild type or dominant active *VKORC1* and binding partners or signaling partners thereof. Compounds are screened for their ability to interact with *VKORC1* proteins and inhibit or enhance *VKORC1* function or the interaction of *VKORC1* with binding partners (e.g., cadherin).

In still further embodiments, cells or transgenic animals having altered (e.g., polymorphic) *VKORC1* genes are utilized in drug screening applications. For example, in some embodiments, compounds are screened for their ability to alter blood clotting in *VKORC1* mice with a particular polymorphic form of *VKORC1*.

In yet other embodiments, subjects (e.g., human subject) are enrolled in clinical trials to test dosages of Warfarin or other related drugs (e.g., new drugs). In preferred embodiments, subjects having polymorphic *VKORC1* are included in clinical trials to test clotting drugs.

The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including biological libraries; peptoid libraries (libraries of molecules having the functionalities of peptides, but with a novel, non-peptide backbone, which are resistant to enzymatic degradation but which nevertheless remain bioactive; see, e.g., Zuckermann *et al.*, J. Med. Chem. 37: 2678-85 [1994]); spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method; and synthetic library methods using affinity chromatography selection. The biological library and peptoid library approaches are preferred for use with peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam (1997) Anticancer Drug Des. 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt *et al.*, Proc. Natl. Acad. Sci. U.S.A. 90:6909 [1993]; Erb *et al.*, Proc. Nad. Acad. Sci. USA 91:11422 [1994]; Zuckermann *et al.*, J. Med. Chem. 37:2678 [1994]; Cho *et al.*, Science 261:1303 [1993]; Carrell *et al.*, Angew. Chem. Int. Ed. Engl. 33:2059 [1994]; Carell *et al.*, Angew. Chem. Int. Ed. Engl. 33:2061 [1994]; and Gallop *et al.*, J. Med. Chem. 37:1233 [1994].

Libraries of compounds may be presented in solution (*e.g.*, Houghten, Biotechniques 13:412-421 [1992]), or on beads (Lam, Nature 354:82-84 [1991]), chips (Fodor, Nature 364:555-556 [1993]), bacteria or spores (U.S. Patent No. 5,223,409; herein incorporated by reference), plasmids (Cull *et al.*, Proc. Nad. Acad. Sci. USA 89:18651869 [1992]) or on phage (Scott and Smith, Science 249:386-390 [1990]; Devlin Science 249:404-406 [1990]; Cwirla *et al.*, Proc. Natl. Acad. Sci. 87:6378-6382 [1990]; Felici, J. Mol. Biol. 222:301 [1991]).

15 IV. Transgenic Animals Expressing *VKORC1* Polymorphic Sequences

The present invention contemplates the generation of transgenic animals comprising an exogenous *VKORC1* gene or mutants and variants thereof (*e.g.*, single nucleotide polymorphisms). In preferred embodiments, the transgenic animal displays an altered phenotype (*e.g.*, response to Warfarin or other anticoagulant drugs) as compared to wild-type animals. Methods for analyzing the presence or absence of such phenotypes include but are not limited to, those disclosed herein.

The transgenic animals or natural variants having equivalent genotypes of the present invention find use in drug (*e.g.*, anticoagulant) screens. In some embodiments, test compounds (*e.g.*, a drug that is suspected of being useful as an anticoagulant therapy) and control compounds (*e.g.*, a placebo) are administered to the transgenic animals and the control animals and the effects evaluated.

The transgenic animals can be generated via a variety of methods. In some embodiments, embryonal cells at various developmental stages are used to introduce transgenes for the production of transgenic animals. Different methods are used depending on the stage of development of the embryonal cell. The zygote is the best target for micro-injection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter that allows reproducible injection of 1-2 picoliters (pl) of DNA solution. The use of zygotes as a target for gene transfer has a major advantage in that in most cases the injected DNA will be incorporated into the host genome before the first

cleavage (Brinster *et al.*, Proc. Natl. Acad. Sci. USA 82:4438-4442 [1985]). As a consequence, all cells of the transgenic non-human animal will carry the incorporated transgene. This will in general also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the transgene.

- 5 U.S. Patent No. 4,873,191 describes a method for the micro-injection of zygotes; the disclosure of this patent is incorporated herein in its entirety.

In other embodiments, retroviral infection is used to introduce transgenes into a non-human animal. In some embodiments, the retroviral vector is utilized to transfect oocytes by injecting the retroviral vector into the perivitelline space of the oocyte (U.S. Pat. No. 10 6,080,912, incorporated herein by reference). In other embodiments, the developing non-human embryo can be cultured *in vitro* to the blastocyst stage. During this time, the blastomeres can be targets for retroviral infection (Janenich, Proc. Natl. Acad. Sci. USA 73:1260 [1976]). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Hogan *et al.*, in *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. [1986]). The viral vector system used to introduce the transgene is typically a replication-defective retrovirus carrying the transgene (Jahner *et al.*, Proc. Natl. Acad. Sci. USA 82:6927 [1985]). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Stewart, *et al.*, EMBO J., 6:383 [1987]). Alternatively, infection 20 can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (Jahner *et al.*, Nature 298:623 [1982]). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of cells that form the transgenic animal. Further, the founder may contain various retroviral insertions of the transgene at different positions in the genome that generally will segregate in the offspring. In addition, 25 it is also possible to introduce transgenes into the germline, albeit with low efficiency, by intrauterine retroviral infection of the midgestation embryo (Jahner *et al.*, *supra* [1982]). Additional means of using retroviruses or retroviral vectors to create transgenic animals known to the art involve the micro-injection of retroviral particles or mitomycin C-treated cells producing retrovirus into the perivitelline space of fertilized eggs or early embryos 30 (PCT International Application WO 90/08832 [1990], and Haskell and Bowen, Mol. Reprod. Dev., 40:386 [1995]).

In other embodiments, the transgene is introduced into embryonic stem cells and the transfected stem cells are utilized to form an embryo. ES cells are obtained by culturing pre-implantation embryos *in vitro* under appropriate conditions (Evans *et al.*, Nature

292:154 [1981]; Bradley *et al.*, Nature 309:255 [1984]; Gossler *et al.*, Proc. Acad. Sci. USA 83:9065 [1986]; and Robertson *et al.*, Nature 322:445 [1986]). Transgenes can be efficiently introduced into the ES cells by DNA transfection by a variety of methods known to the art including calcium phosphate co-precipitation, protoplast or spheroplast fusion, lipofection and DEAE-dextran-mediated transfection. Transgenes may also be introduced into ES cells by retrovirus-mediated transduction or by micro-injection. Such transfected ES cells can thereafter colonize an embryo following their introduction into the blastocoel of a blastocyst-stage embryo and contribute to the germ line of the resulting chimeric animal (for review, *See*, Jaenisch, Science 240:1468 [1988]). Prior to the introduction of transfected ES cells into the blastocoel, the transfected ES cells may be subjected to various selection protocols to enrich for ES cells which have integrated the transgene assuming that the transgene provides a means for such selection. Alternatively, the polymerase chain reaction may be used to screen for ES cells that have integrated the transgene. This technique obviates the need for growth of the transfected ES cells under appropriate selective conditions prior to transfer into the blastocoel.

In still other embodiments, homologous recombination is utilized knock-out gene function or create deletion mutants (*e.g.*, truncation mutants). Methods for homologous recombination are described in U.S. Pat. No. 5,614,396, incorporated herein by reference.

20 EXPERIMENTAL

The following example is provided in order to demonstrate and further illustrate certain preferred embodiments and aspects of the present invention and are not to be construed as limiting the scope thereof.

25 Example 1

***VKORC1* Polymorphisms**

This Example describes the association between *VKORC1* polymorphisms and optimal Warfarin dosages.

30 A. Methods

Clinical and control subjects

The initial European American clinical patients used in this study have been previously described (Higashi *et al.*, Jama 287, 1690-8 (2002)) as have most of the European American patients in the replication study (Gage *et al.*, Thromb Haemost 91, 87-

94 (2004)). All control DNA population samples were purchased from the human variation collections and the CEPH pedigree samples at the Coriell Cell Repository. The Asian American samples consisted of 96 individuals from the HD100CHI panel (Han People of Los Angeles), 10 Southeast Asians (HD13), 7 Chinese (HD32), and 7 Japanese (from the HD07 panel). The 96 European American samples were selected from the HD100CAU panel with the remaining 23 individuals selected from the parental generation of the CEPH families (for more information on these samples see Table 4). The 96 African American samples were selected from the HD100AA panel.

10 Sequence analysis and genotyping

All clinical samples from the primary European American cohort were resequenced for SNP discovery using PCR amplification of ~1 kb fragments covering the entire genomic region of *VKORC1* and direct sequencing of the PCR amplicons using standard ABI Big-Dye Terminator sequencing chemistry and run on an ABI 3730XL DNA analyzer. SNPs were identified using the program Polyphred (v. 4.2), along with quality control and review of all SNPs and genotypes by a human analyst. The ten SNPs identified were at position 381(C/T), 861(A/C), 2653(G/C), 3673(A/G), 5808(T/G), 6009(C/T), 6484(C/T), 6853(C/G), 7566(C/T), and 9041(A/G) in the *VKORC1* reference sequence (GenBank Accession AY587020; SEQ ID NO:1). A single heterozygous non-synonymous SNP was identified (genomic position 5432 (G/T) – Ala41Ser) in a European American clinical patient. This patient had the highest overall warfarin maintenance dose (15.5 mg/d) and was excluded from all analyses. No other previously reported nonsynonymous SNPs were identified (Rost et al., Nature 427, 537-41 (2004)). All other control population samples were resequenced using the same methods, but genotyped using only the amplicons containing the 10 common SNPs identified in the European American clinical population.

For the replication study in the secondary European American cohort, four informative SNPs (861, 5808, 6853, and 9041) were used to differentiate between haplotype H1, H2, H7, H8 and H9, based on the genealogical tree in Fig. 1. For each SNP site, PCR primers were designed using Primer Express version 1.5 (ABI, Foster City, CA). Pyrosequencing primers were designed using the Pyrosequencing SNP Primer Design Version 1.01 software. Unique localization of the PCR primers was verified using NCBI Blast (available at the Internet site of NCBI). PCR was carried out using Amplitaq Gold PCR master mix (ABI, Foster City, CA), 5 pmole of each primer (IDT, Coralville, IA), and 1ng DNA. Pyrosequencing was carried out as previously described (Rose et al., Methods

Mol Med 85, 225-37 (2003) using the following primers (5' – 3') for each SNP: 861 (A/C), forward = TCTTGGAGTGAGGAAGGCAAT (SEQ ID NO:2), reverse = Biotin-GACAGGTCTGGACAACGTGG (SEQ ID NO:3), internal = CTCAGGTGATCCA (SEQ ID NO:4); 5808 (G/T), forward = Biotin - GGATGCCAGATGATTATTCTGGAGT (SEQ ID NO:5), reverse = TCATTATGCTAACGCCTGGCC (SEQ ID NO:6), internal = CAACACCCCCCTTC (SEQ ID NO:7); 6853 (G/C), forward = CTTGGTGATCCACACAGCTGA (SEQ ID NO:8), reverse = Biotin - AAAAGACTCCTGTAGTTACCTCCCC (SEQ ID NO:9), internal = AGCTAGCTGCTCATCAC (SEQ ID NO:10); 9041 (A/G), forward = TACCCCTCCTCCTGCCATA (SEQ ID NO:11), reverse = Biotin - CCAGCAGGCCCTCCACTC (SEQ ID NO:12), internal = TCCTCCTGCCATAACC (SEQ ID NO:13). Samples of each genotype were randomly selected and repeated to confirm the genotype assignment.

15 Statistical methods

Genealogic trees were constructed using the program MEGA and based on the number of differences between haplotypes and the UPGMA clustering method. Haplotypes for each individual sample were estimated using the program PHASE, version 2.0 (Stephens and Donnelly, Am J Hum Genet 73, 1162-9 (2003)), and independent runs were performed for each population studied.

Using the most likely pair of haplotypes estimated for each patient, the association between number of copies of each *VKORC1* haplotype (coded 0, 1, 2) and maintenance warfarin dose was assessed on an additive scale. Multiple linear regression was performed using log-transformed maintenance dose, adjusting for the covariates age, sex, race, amiodarone, losartan, and *CYP2C9* genotype. Adjusted warfarin doses (and 95% confidence intervals) associated with each additional haplotype copy were estimated by exponentiation of the mean fitted values and standard errors of the linear prediction. In separate analyses, using a generalized linear model score test method (Lake et al., Hum Hered 55, 56-65 (2003)) that additionally takes into account the uncertainty of haplotypes assignments, similar estimates were obtained for mean warfarin dose, and the confidence values were slightly wider.

The Kruskal-Wallis test, a distribution-free ANOVA, was used to assess differences in maintenance dose among the A/A, A/B and B/B groups. This was done separately for three subsets of the data: (1) for subjects with the *2 or *3 variant, (2) wild type and (3) *2

or *3 and wild type combined. Subjects with a non-A or B haplotype were not used in the analysis. Following the overall chi-square test for differences among the three groups, pairwise comparisons of groups were carried out using the asymptotic normality of the total ranks within each group. The Bonferroni correction for each of the three individual
5 comparisons (A/A vs A/B, A/B vs B/B, and A/A vs B/B) was made to control the overall type I error rate.

Differences between population specific haplotype distributions were done using a χ^2 test.

10 B. Results

In order to investigate the link between common, non-coding single nucleotide polymorphisms (SNPs) in *VKORC1* and warfarin dosing, complete gene resequencing of the *VKORC1* gene locus (11.2 kilobases) in a cohort of 185 European American patients receiving long-term warfarin therapy was carried out. All patients had been previously
15 genotyped for known functional *CYP2C9* mutations (*2 and *3) that are associated with lower warfarin dose requirements (Higashi et al., *Jama* 287, 1690-8 (2002); Aithal et al., *Lancet* 353, 717-9 (1999)). In *VKORC1*, all clinical samples were resequenced over 5 kilobases in the upstream promoter region, 4.2 kilobases of intragenic (intron and exon) sequence, and 2 kilobases of the 3' downstream region. Ten non-coding SNPs with a minor
20 allele frequency greater than 5% were identified in the European American clinical patients. These SNPs were used to estimate *VKORC1* haplotypes that were assigned to each patient. From these 185 patients, five common haplotypes (>5%) were identified – H1, H2, H7, H8, H9 (Table 1).

When each SNP was tested individually, seven sites (381, 3673, 5808, 6484, 6853,
25 7566, and 9041) were highly significant ($p < 0.001$) and three sites were marginally significant (861, 2653, and 6009, $p = 0.01$, 0.02, and 0.02, respectively) when regressed against daily warfarin maintenance dose. Of the seven highly significant sites, five (381, 3673, 6484, 6853, 7566) are in strong linkage disequilibrium ($r^2 = 0.9$) and two independent sites (5808 and 9041) are not correlated with any other SNP in this region. Analysis of
30 SNP-SNP interactions also showed significant effects between multiple site combinations, therefore, the association of individual haplotypes with warfarin doses was also quantified. A multiple linear regression analysis using inferred haplotypes for each patient was used to determine the association of haplotype on warfarin dose, while adjusting for genetic and other clinically important covariates (e.g. age, *CYP2C9*-*2 or *3, etc; see Table 1 and Table

4). Four of the five common haplotypes (frequency > 5%) were found to be significantly associated with warfarin dose ($p \leq 0.05$) (Table 1). From this analysis, two low-dose (2.9 and 3.0 mg/d) haplotypes (H1 and H2) and two high-dose (6.0 and 5.5 mg/d) haplotypes (H7 and H9) were identified.

5 A genealogical tree was constructed from the five common haplotypes to identify potential hierarchical haplotype groupings (Fig. 1 – upper panel). Two distinct haplotype clades, which were completely segregating at five of the ten *VKORC1* SNPs, were identified and designated clade A (H1 and H2) and clade B (H7, H8, and H9). Using this designation, all patients were grouped based on their *CYP2C9* genotype and assigned a *VKORC1* clade
10 diplotype (i.e. combination of two clades) of A/A, A/B, or B/B. Fig. 1 – lower panel). The overall mean (5.1 ± 0.2 mg/d) and range of warfarin maintenance doses were typical of other studies of clinical patients (Aithal et al., supra). Warfarin maintenance dose differed significantly between all three clade diplotype groupings (A/A, A/B, B/B, $p < 0.001$) in the combined patient set (i.e. Fig. 1 – All patients), and for the *CYP2C9* wild-type (WT)
15 patients--there was an additive effect over the entire warfarin dose range. Overall, the proportion of warfarin dose variance explained by *VKORC1* clades A and B was 25%, and was similar to values obtained when considering all *VKORC1* SNP sites with interactions. Patients who were carriers of *CYP2C9* *2 or *3 mutations showed a similar effect of *VKORC1* clade diplotype on warfarin dose ($p < 0.001$ between diplotype A/A and A/B).
20 There was an overall trend towards lower warfarin dose associated with *CYP2C9* variant genotype (Fig 1, lower panel), consistent with the known blunted metabolism of warfarin in carriers of these allelic variants (Rettie et al., *Epilepsy Res* 35, 253-5 (1999)). The segregation of *VKORC1* haplotypes into low and high dose associated clades, independently of *CYP2C9**2 and *3, suggests that *VKORC1* SNP genotyping have strong predictive power
25 for determining the warfarin dose needed to achieve and maintain therapeutic anticoagulation in the clinical setting.

In order to validate these initial results, a replication study was performed in a larger, independent cohort of warfarin-treated European American patients ($n = 368$). These patients were genotyped using four informative SNPs (861, 5808, 6853, 9041 – Fig. 1
30 – upper panel – bold numbers) that resolve all five common haplotypes (H1, H2, H7, H8, and H9) present in the initial European American clinical cohort. Haplotypes were inferred, clade diplotypes assigned, and patients segregated based on their known *CYP2C9* genotype. Overall, the results from this larger clinical population recapitulated the salient findings in the index population for all three patient subgroups. In this second cohort, the *CYP2C9*-WT

patients ($n = 233$) and all patients ($n = 357$) showed a significant additive effect across the A/A (3.4 ± 0.26 and 3.2 ± 0.21 mg/d), A/B (4.9 ± 0.17 and 4.4 ± 0.13 mg/d) and B/B (6.7 ± 0.29 and 6.1 ± 0.23 mg/d) clade diplotypes ($p < 0.05$ between A/A and A/B, A/B and B/B).

One variable used in estimating clinical warfarin dose is racial background of the patient (Blann et al., Br J Haematol 107, 207-9 (1999); Gan et al., Int J Hematol 78, 84-6 (2003)). Individuals of Asian-, European-, and African ancestry tend to require, on average, lower (~ 3.0 mg/d), intermediate (~ 5.0 mg/d) and higher (~ 6.5 mg/d) dose, respectively (Yu et al., Qjm 89, 127-35 (1996); Chenhsu et al., Ann Pharmacother 34, 1395-401 (2000); Absher et al., Ann Pharmacother 36, 1512-7 (2002); Gage et al., Thromb Haemost 91, 87-94 (2004)). In order to investigate whether this variation in dose requirement may be due to population specific differences in the distribution of *VKORC1* haplotypes, 335 unrelated control individuals, selected from these population ancestries (European, $n = 119$, African, $n = 96$, Asian, $n = 120$) were resequenced and the genotype was determined at each of the 10 SNPs present in the European-descent clinical patients. Haplotype pairs for each individual were inferred, and the population haplotype frequencies determined along with the distribution of clade A and B haplotypes (Table 2). The distribution of common haplotypes (H1, H2, H7, H8, and H9) between the European American clinical and control populations was significantly different ($p < 0.001$), primarily due to an increase in the high dose associated H7 haplotype in clinical patients. This may be due to selection bias in the clinical population resulting from preferential referral, to an academic medical center, from which the patients were recruited.

The five predictive haplotypes accounted for 99% and 96%, of the total haplotypes in the European American clinical and control populations; no significant difference was present based on the distribution of clades A (35% vs 37%) and B (64% vs 58%). The five common haplotypes within the European American population accounted for only 61% of total African American haplotypes. This more diverse distribution of haplotypes in the African American population is consistent with the higher genomic sequence diversity found in African-descent populations (Przeworski et al., Trends Genet 16, 296-302 (2000); Crawford et al., Am J Hum Genet 74, 610-22 (2004)). These population-specific haplotype differences may be due to demographic effects such geographic selective pressures, migration, or bottlenecks, and have been observed for other medically relevant genes (e.g. *ADRB2*, (Drysdale et al., Proc Natl Acad Sci U S A 97, 10483-8 (2000)). The African and Asian American populations showed significant differences in clade A and B frequencies (p

< 0.001) compared to the European American control population. The frequency of clade A haplotypes was higher among the Asian American population (89%) and lower in the African American population (14%) compared to the European American control population (37%). Because clade A haplotypes predict the low warfarin dose phenotype (Table 1), ethnic differences in *VKORC1* haplotype frequency parallel the clinical experience of population differences in warfarin maintenance dose requirements. Thus, this example describes population specific differences in haplotype distribution that are a major contributor to the variation in warfarin maintenance dose requirements between racial groups.

The molecular mechanism(s) by which these haplotypes, or the individual SNP alleles that comprise them, determine warfarin dose remain undefined. Two of these SNPs (381 and 3673) are present in the 5' upstream promoter region, two in the first intron (5808 and 6484) and one (9041) in the 3' untranslated region (UTR). None of the significantly associated SNPs are present in highly conserved non-coding sequence present in mouse or rat. The present invention is not limited to a particular mechanism. Indeed, an understanding of the mechanism is not necessary to practice the present invention. Nonetheless, it is contemplated that SNPs in the 3' UTR may affect mRNA folding and stability (Durrin, L. K., Haile, R. W., Ingles, S. A. & Coetzee, G. A. Vitamin D receptor 3'-untranslated region polymorphisms: lack of effect on mRNA stability. *Biochim Biophys Acta* 1453, 311-20 (1999); Carter, A. M., Sachchithanathan, M., Stasinopoulos, S., Maurer, F. & Medcalf, R. L. Prothrombin G20210A is a bifunctional gene polymorphism. *Thromb Haemost* 87, 846-53 (2002)), which could alter *VKORC1* expression and possibly warfarin response. The present invention is not limited to a particular mechanism. Indeed, an understanding of the mechanism is not necessary to practice the present invention. Nonetheless, it is contemplated that the strong association of individual haplotypes with warfarin dose also suggests that a functional interaction between SNP alleles carried on the same haplotype may be contributing to the observed results.

In summary, this Example describes *VKORC1* noncoding SNPs and haplotypes that are strongly associated with warfarin dose. These haplotypes group into higher order clades that segregate patients into low, intermediate and high warfarin maintenance doses. The *VKORC1* gene-warfarin dose association is independent of *CYP2C9* genotype, and explains 23%–25% of the variability in the warfarin dose. Genotyping for these *VKORC1* SNPs and haplotypes provides more accurate initial dosing and reduces the amount of time to stable

anticoagulation, thereby improving the safety, effectiveness, and hospitalization costs associated with warfarin therapy.

5

Table 1. Average warfarin maintenance dose requirement based on *VKORC1* haplotype.

Haplotype	Haplotype Sequence	Frequency	Average maintenance dose for homozygous patients (mg/d)*
H1	CCGATCTCTG (SEQ ID NO:14)	0.12	2.9 (2.2 – 3.7)
H2	CCGAGCTCTG (SEQ ID NO:15)	0.24	3.0 (2.5 – 3.6)
H7	TCGGTCCGCA (SEQ ID NO:16)	0.35	6.0 (5.2 – 6.9)
H8	TAGGTCCGCA (SEQ ID NO:17)	0.08	4.8 (3.4 – 6.7)
H9	TACGTTCGCG (SEQ ID NO:18)	0.21	5.5 (4.5 – 6.7)

10 *Adjusted for race, age, sex, amiodarone, losartan, and CYP2C9 variant genotype. Warfarin dose effect for each haplotype is shown as: Mean (95% confidence interval). p-values for each haplotype were H1, $p < 0.0001$; H2, $p < 0.001$; H7, $p < 0.001$; H8, $p = 0.76$, and H9 $p = 0.05$. $n = 185$ clinical samples.

15 Note: For each haplotype sequence the alleles are listed in sequential order across the *VKORC1* gene – 381, 861, 2653, 3673, 5808, 6009, 6484, 6853, 7566, and 9041.

Table 2. *VKORC1* haplotype distributions in European-, African- and Asian American populations.

Haplotype	Haplotype Sequence	European Clinic	European Controls	African Controls	Asian Controls
1	CCGATCTCTG (SEQ ID NO:14)	43 (0.12)	28 (0.12)	14 (0.07)	213 (0.89)
2	CCGAGCTCTG (SEQ ID NO:15)	88 (0.24)	61 (0.26)	12 (0.06)	0 (0.00)
3	CCGGTCCCCG (SEQ ID NO:19)	2 (0.01)	3 (0.01)	27 (0.14)	0 (0.00)
4	CCGGTCCGTG (SEQ ID NO:20)	1 (0.00)	0 (0.00)	11 (0.06)	0 (0.00)
5	TCGAGCTCTG (SEQ ID NO:21)	1 (0.00)	5 (0.02)	0 (0.00)	0 (0.00)

	TCGGTCCGCG				
6	(SEQ ID NO:22)	0 (0.00)	0 (0.00)	15 (0.08)	0 (0.00)
	TCGGTCCGCA				
7	(SEQ ID NO:16)	132 (0.35)	49 (0.21)	80 (0.42)	25 (0.10)
	TAGGTCCGCA				
8	(SEQ ID NO:17)	28 (0.08)	34 (0.14)	2 (0.01)	0 (0.00)
	TACGTTCGCG				
9	(SEQ ID NO:18)	77 (0.21)	56 (0.24)	11 (0.06)	0 (0.00)
OTHER	-	0 (0.00)	2 (0.01)	20 (0.10)	2 (0.01)
	Clade A (1,2)	131 (0.35)	89 (0.37)	26 (0.14)	213 (0.89)
	Clade B (7,8,9)	237 (0.64)	139 (0.58)	93 (0.47)	25 (0.10)
	TOTAL(A&B)	340 (0.99)	194 (0.96)	119 (0.61)	238 (0.99)
Total Chromosomes (2N)		372	238	192	240
Total Individuals (N)		186	119	96	120

Note: Haplotype alleles at each position are listed in the same order as Table 1. For each population the number of inferred haplotypes is listed. Numbers in parentheses denote proportion of individuals with given haplotype.

5

Table 3. *VKORC1* SNP genotype tests and maintenance warfarin dose

<i>Genotype</i>	<i># (%)</i>	<i>Mean dose (95% CI)</i>	<i>Unadjusted P-value</i>	<i>Adjusted P-value*</i>
<i>VKORC1</i> 381			<0.001	<0.001
C/C	49 (42)	5.4 (4.7 – 6.3)		
C/T	56 (47)	4.6 (4.2 – 5.1)		
T/T	13 (11)	2.3 (1.8 – 2.9)		
<i>VKORC1</i> 861			0.01	0.01
A/A	58 (48)	4.0 (3.5 – 4.5)		
A/C	49 (40)	5.0 (4.4 – 5.7)		
C/C	14 (12)	5.3 (4.2 – 6.6)		
<i>VKORC1</i> 2653			0.009	0.02
G/G	115 (64)	4.3 (3.9 – 4.7)		
G/C	59 (33)	4.9 (4.3 – 5.6)		
C/C	7 (4)	6.6 (4.3 – 10.2)		
<i>VKORC1</i> 3673			<0.001	<0.001
A/A	77 (43)	5.5 (4.9 – 6.2)	<0.001	<0.001
G/A	81 (45)	4.6 (4.2 – 5.0)	0.004	0.005
G/G	22 (12)	2.6 (2.2 – 3.1)	<0.001	<0.001
<i>VKORC1</i> 5808			<0.001	0.0001
T/T	104 (60)	5.2 (4.8 – 5.7)		
T/G	60 (35)	4.0 (3.6 – 4.6)		
G/G	9 (5)	2.6 (2.0 – 3.5)		

<i>VKORC1</i> 6009			0.007	0.02
C/C	110 (62)	4.3 (3.9 – 4.7)		
C/T	61 (34)	5.0 (4.4 – 5.6)		
T/T	7 (4)	6.6 (4.3 – 10.2)		
<i>VKORC1</i> 6484			<0.001	<0.001
C/C	77 (42)	5.5 (4.9 – 6.2)		
C/T	83 (46)	4.5 (4.1 – 4.9)		
T/T	22 (12)	2.6 (2.2 – 3.1)		
<i>VKORC1</i> 6853			<0.001	<0.001
C/C	72 (41)	5.5 (4.8 – 6.1)		
C/G	80 (46)	4.4 (4.1 – 4.8)		
G/G	22 (13)	2.6 (2.2 – 3.0)		
<i>VKORC1</i> 7566			<0.001	<0.001
C/C	74 (42)	5.4 (4.8 – 6.1)		
C/T	83 (47)	4.5 (4.1 – 4.9)		
T/T	21 (12)	2.6 (2.2 – 3.0)		
<i>VKORC1</i> 9041			<0.001	<0.001
A/A	56 (32)	3.7 (3.3 – 4.3)		
G/A	87 (50)	4.8 (4.4 – 5.3)		
G/G	30 (17)	5.9 (5.2 – 6.6)		

*Adjusted for age, race, sex, amiodarone, losartan, CYP*2, and CYP*3.

P-values were derived from likelihood ratio test statistics of linear regression models in which the number of SNP alleles was coded 0,1, 2 to represent an additive or co-dominant genetic model of inheritance.

5

Table 4. Characteristics of 185 European American Warfarin Clinic Patients

Characteristic	N (%) or mean ± SD (range)
Sex	
Male	121 (65)
Female	64 (35)
Race	
White	179 (97)
Hispanic	6 (3)
Age, years (range)	59.9 ± 15.7 (19 – 88)
Cigarette smoker	25 (14)
Diagnosis	
Atrial fibrillation	95 (52)

Arrhythmia	81 (44)
Congestive heart failure	77 (42)
Venous thromboembolic disease	40 (22)
Dilated cardiomyopathy	37 (20)
Valvular disease	13 (7)
Hypertension	85 (46)
Diabetes type	41 (22)
Malignancy	27 (15)
Medication use	
Amiodarone	24 (13)
Losartan	17 (9)
Torsemide	11 (6)
Acetaminophen	52 (28)
Vitamin C	27 (15)
Vitamin E	25 (14)
Maintenance warfarin dose, mg/day	5.1 ± 2.5
Follow-up, days	
Mean	831
Median	545
Range	14 - 4032

5 **Table 5. Comparison of daily warfarin dose and clade diplotype between the two European American clinical cohorts.**

Index Population (n = 185) – Seattle – University of Washington

	AA			AB			BB		
	ALL	*2 or *3	WT	ALL	*2 or *3	WT	ALL	*2 or *3	WT
Average	2.58	2.37	2.69	4.79	4.00	5.15	6.23	4.40	7.00
StDev	0.82	0.87	0.79	1.83	1.10	1.98	2.71	1.41	2.77
SEM	0.17	0.31	0.20	0.20	0.21	0.26	0.32	0.30	0.38
n	23	8	15	86	27	59	74	22	52

Replication Population (n = 368) – St. Louis - Washington University

	AA			AB			BB		
	ALL	*2 or *3	WT	ALL	*2 or *3	WT	ALL	*2 or *3	WT
Average	3.20	2.78	3.35	4.42	3.61	4.90	6.11	5.00	6.68
StDev	1.40	1.21	1.46	1.75	1.40	1.77	2.71	2.08	2.83
SEM	0.21	0.35	0.26	0.13	0.18	0.17	0.23	0.30	0.29
n	44	12	32	170	63	107	143	49	94

Note: some individuals were not able to be classified within the A or B clades

Example 2**Expression of *VKORC1* mRNA**

To explore the mechanism of the association between warfarin dose and *VKORC1* polymorphisms, *VKORC1* mRNA levels were assayed in human liver tissue and compared with the major *VKORC1* haplotype groups (A/A, A/B, B/B).

A. Methods

Assay of VKORC1 mRNA. 1.2 μ L of total cDNA from each sample was used as template for the quantitative PCR (using 9 μ L reactions) in the presence of SYBR green reporter (Applied Biosystems, Foster City, CA). PCR primers (5' to 3' - forward = ATCAGCTGTTCGCGCGTC (SEQ ID NO:14), reverse = AGAGCACGAAGAACAGGATC (SEQ ID NO:15) were selected from sequences in exon 1 and 3 of the *VKORC1* coding sequence (Accession No. NM_024006). All quantitative PCR was performed on an Applied Biosystems 7900HT and real-time data collected during the entire thermocycling period (cycling conditions: 95°C – 15 minutes for initial denaturation and 40 cycles of 94°C – 30 sec., 60 °C – 30 sec, 72°C – 30 sec and a final extension of 72 °C – 5 minutes). Each sample was measured in duplicate and the results from two independent experiments were averaged. All *VKORC1* mRNA levels were normalized to GAPDH expression levels (primers (5' to 3'): forward = ACAGTCAGCCGCATCTTCTT (SEQ ID NO:16), reverse = ATGGGTGGAATCATATTGGAAC (SEQ ID NO:17), and scaled relative to the A/A haplotype group (Mean value = 1.49).

Liver mRNA expression data were analyzed following log transformation and the overall test for group differences was performed using an ANOVA. Pairwise comparisons between groups for significance were performed using Tukey's Studentized Range Test. Significance levels were set at $p < 0.05$.

B. Results

Results are shown in Figure 3. A graded and highly significant ($p = 0.002$) gene-dose effect is evident, with mRNA levels about 3-fold higher in the B/B ('high-dose') group compared to the A/A ('low-dose' group) (Figure 3 - $p < 0.05$).

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention that are obvious to those skilled in the art are intended to be within the scope of the following claims.

10

15

CLAIMS

We claim:

1. A method, comprising: providing a sample from a subject; and determining
5 said subject's *VKORC1* haplotype to determine responsiveness to Warfarin therapy.
2. The method of claim 1, further comprising the step of determining said
subject's optimal Warfarin dose based on said subject's *VKORC1* haplotype.
- 10 3. The method of claim 1, wherein said subject's haplotype is selected from the
group consisting of H1, H2, H3, H4, H5, H6, H7, H8, and H9.
4. The method of claim 1, further comprising the step of determining said
subject's CYP2C9 genotype.
15
5. The method of claim 1, wherein said determining said subject's *VKORC1*
haplotype comprises a nucleic acid based detection assay.
6. The method of claim 5, wherein said nucleic acid based detection assay is
20 selected from the group consisting of a sequencing assay and hybridization assay.
7. The method of claim 1, further comprising the step of determining said
subject's Clade type.
- 25 8. The method of claim 7, wherein said Clade type is selected from the group
consisting of AA, AB, and BB.

9. A method, comprising:
- a) providing a sample from a subject;
 - b) detecting the genotype of a single nucleotide polymorphism at one or more positions selected from the group consisting of positions 381, 3673, 5808, 6484, 6853, 7566, and 9041 of SEQ ID NO:1 and positions in linkage disequilibrium with said positions; and
 - c) determining said subjects optimal Warfarin dosage based on said genotype of said single nucleotide polymorphism.
10. The method of claim 9, further comprising the step of determining said subject's CYP2C9 genotype.
11. The method of claim 9, wherein said determining said subject's *VKORC1* haplotype comprises a nucleic acid based detection assay.
12. The method of claim 11, wherein said nucleic acid based detection assay is selected from the group consisting of a sequencing assay and hybridization assay.
13. A kit for determining a subject optimal Warfarin dosage, comprising:
- a) a detection assay, wherein said detection assay is configured to specifically detect said subject's *VKORC1* haplotype; and
 - b) instructions for determining said subject's optimal Warfarin dosage.
14. The kit of claim 13, wherein said subject's *VKORC1* haplotype is selected from the group consisting of H1, H2, H7, H8, and H9.
15. The kit of claim 13, further comprising reagents for determining said subject's CYP2C9 genotype.
16. The kit of claim 13, wherein said detection assay is a nucleic acid based detection assay.
17. The kit of claim 16, wherein said nucleic acid based detection assay is selected from the group consisting of a sequencing assay and hybridization assay.

18. The kit of claim 13, wherein said kit further comprises instructions for determining said subject's Clade type.

5 19. The kit of claim 18, wherein said Clade type is selected from the group consisting of AA, AB, and BB.

20. A method, comprising: providing a sample from a subject; and determining said subject's *VKORC1* expression level to determine responsiveness to Warfarin therapy.

10

21. The method of claim 20, further comprising the step of determining said subject's optimal Warfarin dose based on said subject's *VKORC1* expression level.

22. The method of claim 20, further comprising the step of determining said
15 subject's CYP2C9 genotype.

23. The method of claim 20, wherein said determining said subject's *VKORC1* expression level comprises determining the amount of *VKORC1* mRNA expressed by said subject.

20

24. The method of claim 23, wherein said determining the amount of *VKORC1* mRNA expressed by said subject comprises a quantitative RT-PCR assay.

25. The method of claim 23, wherein said determining the amount of *VKORC1*
25 mRNA expressed by said subject comprises a nucleic acid hybridization assay.

26. The method of claim 20, wherein said determining said subject's *VKORC1* expression level comprises determining the amount of *VKORC1* polypeptide expressed by said subject.

30

27. The method of claim 26, wherein said determining the amount of *VKORC1* polypeptide expressed by said subject comprises exposing said sample to an antibody that specifically binds to said *VKORC1* polypeptide.

28. A kit for determining a subject optimal Warfarin dosage, comprising:

a) reagents for performing a detection assay, wherein said detection assay is configured to specifically detect said subject's *VKORC1* expression level; and

5 b) instructions for determining said subject's optimal Warfarin dosage.

29. The kit of claim 28, wherein said reagents comprise reagents for determining the amount of *VKORC1* mRNA expressed by said subject.

10 30. The kit of claim 29, wherein said reagents comprise reagents for performing a quantitative RT-PCR assay.

31. The kit of claim 29, wherein said reagents comprise reagents for performing a nucleic acid hybridization assay.

15

32. The kit of claim 28, wherein said reagents comprise reagents for determining the amount of *VKORC1* polypeptide expressed by said subject.

20 33. The kit of claim 32, wherein said reagents comprise an antibody that specifically binds to said *VKORC1* polypeptide.

34. The kit of claim 28, further comprising reagents for determining said subject's CYP2C9 genotype.

Figure 1




Figure 2
SEQ ID NO: 1

```

1   cagaggacaa catagatagg ttcggagact actcagagaa agggagagag aatgctaggt
61  tatctttcag ctgttttagc tgcttattgt tccagcttta gccaccatct gactgcaatt
121 gtatgagcca gaaatagcca ggtaaaattg tcttcccaaa ttcttagtgc acaaaaactg
181 tcagaggctg ggggcagtg ctcactcctg taatcccagc agcactttgg gaagccaggg
241 tgggagaact gcttgagccc aggagttaga ggcaacataa caacaactca tctctaataa
301 aaaaaaaaaa aaaagaaaaga aaaaggccag gcacagtggc tcacgcctgt aatcccagca
361 ctttagggaag ccaaggaggg tggatcatga ggtcaggagt ttgagaccag cctggccaat
421 atggtgaaac cccgtctcta ctaaaaacac aaaaatcagc cagcatagt ggctggtgcc
481 tgtaatcaca gctactcagg aggctaaggc aggagaatca cttgaatccg ggaggcagag
541 gttgcagtga gccgagattg caccactgca tttcagcctg ggcagcagag tgggactgtc
601 tcacattaat taattaattg attaaatgct agaagaaaat atggtcaaaa tcttttataa
661 ctcttgaggt gaggaaggca atctcggtc actacaactt ccacctcccg ggttcaaatg
721 attctcctgc ctcagcctcc tgagtagctg ggattacagg caccaccac catgcccggc
781 taatttttgt atttttagta gagacgaggt ttcgccatgt tggccaagct tgtcttaaac
841 tctgacctc aggtgatcca cccacctcag cctcccaaat tgctgggatt acaggcatga
901 gccaccatgc cccgcctaatt tttaaaaatg tttataaaga cagagtctta ccacgtgtgc
961 cagacctgtc tcaaacctct gggctcaagt cattgtcttg tctcagctc ccaaagtctt
1021 gggactacag gtgtgcacca ccacacctgg ctaattttgt ttatttttta tagagttagg
1081 gtctccctgt gtgtccagg gtgtctcaaa ctctgggct caagtggctc tctgctca
1141 gcctcccaaa gtgtgggat tataggcata agccactgca cctggcccca gtgtgtgtct
1201 tctgaggcta agtcataaag agattgcggc ttctattttg cttcctctta gatcactcat
1261 tctgggggaa accaacaata aataacagga acacaccagg agatcagaga aagctgagat
1321 acaggcctaa tgcccagtc tctgtattca tttctaattg ctgctgtaac aagttactgc
1381 acatttaatg atgtaaaaca acacaaatth atcttacagt tctgtagctt agaagctga
1441 cagagtctca ctgggctgaa atcaagatgt tgatagtctc ttcccgggta ctctagagtg
1501 agaatccagt tctttaccct tcttgctct tagaggccac ccgtatttat tagctcacia
1561 tccccttctt ccatcttcaa accagaacaa ttgcagctct ctgtgtcttt tttctttact
1621 ccatctccc tctgactttc tcttgccatc ctcttcatt ttttaaggac cctgtgggc
1681 caggcatggt ggcttatgcc tgtaatccca gcactttgag aggcggaggc aggcgggtca
1741 cctgagggtc ggagttccag accagcctgg ccaacatggc aaaaccccat ctccactgaa
1801 aatacaaaaa ttggggccagg cagggcggt caccaccgta atccccatac tttgagaggc
1861 tgaggcaggt ggatcacttg aggtcaggag ttcaagacca gctgggcaa catggtgaaa
1921 tcccgctctc actaaaaaaa aaaaaaatta caataattag tcaggcgtgg tggccagcac
1981 ctataatccc agctacttgg gaggttgagg cacaagaatt gcttgaacct gagaggggga
2041 gggtgcagtg agctgagatt gcaccaccgt actccagcct ggatgacaga gcaagactct
2101 gtctcaaaat aaaaattaag ataaataata aaaaaataat aaaaatcagc aaattaggcc
2161 aggtgtgggt gtggctcatg cctgtactcc cagcactttg aggtgggaag gttgcttgaa
2221 accagaagtt tagaccagcc tgggaacaa agtgagacc catctctaca aaaataaaaa
2281 taaatttagcc aggtgtgggt ggacgtactt gtactcctag ctactaggga ggctgaggtg
2341 caagaatcgc ttgaacctgg gagacggagg ttgcagttag ttatgacagt gccactacac
2401 tccagcctgg gtgtcaagg tctttgaaaa aataaacaata ataggctggg cgtggtggct
2461 tacacttgta atcccagcac tttgggaggc tgaggcaggc agatcacaaa gtcaggagtt
2521 cgagaccagc ctggccaaca cagtgaacc ccgtctctac taaaaataca aaaattagcc
2581 aggcaggtt gcacgcgcct gtagtcccaa ctgctcgga ggctgaggca ggagaatggc
2641 ttgaacctgg gggggggcgg aagttataat gagccaagat tgtgccactg cactccagcc
2701 taggcaacag agcgagactc cgtctcaaaa ataaaaataa aaataaatca atagacctg
2761 gtatgatggc tcacgcctat aatcccagca ctttgggagg ccaggtggg tggatcatct
2821 gaggtcagga gtttgagacc agcctgacca acatggagaa accccatctc tactaaaaat
2881 acaaaaaatt agccaggcgt ggtggcacat gcctataatc ccagctactc aggagcctga
2941 ggcaggagaa tcgctttaac cggggaggtg gaggttgagg tgagtcgaga tagcaccatt
3001 gcattctagc ctgggcaaca agagcaaac tccatctcaa aaataaataa ataaatagat

```



```

3061 agataggctg ggcacggtgg ctcacgcctg taatcccagc actttgggag gccgagggtg
3121 gcggatcacc tgaggtcagg agttcaagac cagcctggcc aatatagcga aacctgtgtc
3181 ctactaaaaa tttaaaaaat tagccaggca tgatggcggg tgcctgtaat ccagctactc
3241 cgggaggctg aggcaggaga atctctcgaa cctggcaggc ggaggttgca gtgagccgag
3301 atcacacccat tgcactccag ttgggcaaca agagcgaaac tccacctcaa aaacaaaata
3361 aaataaaata aaatcaggag attggtcagc ttaattccat ctgcaacctt aattcccttt
3421 tggccatttta acctatcata ttcacaagtt ccagggattc atgcaggggac atctttggtg
3481 accattatct tgtctaccac actctctaga agagagtaga tgtgagaaac agcatctgga
3541 gagggaggag ccagcaggag agggaaatat cacagacgcc agaggaaagag agttcccgag
3601 agggtaggtg caacagtaag ggatccctct gggaagtcaa gcaagagaag acctgaaaaa
3661 caaccattgg ccgggtgctg tggctcacgc ctataatcct agcattttgg gaggccgagg
3721 tgggtggatc acttgaggtc aggagtttaa gacaagcctg gccaacatgg tgaaaccctg
3781 tctctactaa aaatacaaaa attagccaga catggtggca ggcacctgta gtccaaacta
3841 cttgggagggc tgaggcacga gaaccacttg aacctgggag gcggaagttg cagttagccg
3901 agatggcacc gctgcactcc agcctgggag acagagttag actccgtctc aaaataagta
3961 aataaataaaa taaataacca ttgggtctag cgtcttggtg acaccagcca ctagggtgtc
4021 gggcaatgct ggaggcagca gctagtttat ggcgtgggca gcaaggaga agtgaggaag
4081 tagagactgg gagtgtagaa tctttccaaa aacttgaatg tggaggggta gatgggataa
4141 tgccgggaga ggaggctggt gaaggacctc ttagtgaagg gagatttgtt gctgtttctg
4201 agatgggaaa gagaacaacg ttgcagtgtg gatggggagg atggagaggt tgaagggtgt
4261 ggagagatgg gtcaccatca gatgggacgt ctgtgaagga gagacctcat ctggcccaac
4321 gcttgggaaag gagagactga ctgttgagtt gatgcaagct caggtgttgc caggcgggag
4381 ccatgatagt agagaggtta ggatactgtc aagggtgtgt gtggccaaag gagtgttctc
4441 gtgaatgtat gggagaaagg gagaccgacc accaggaagc actggtgagg caggaccggg
4501 gaggatggga ggctgcagcc cgaatggtgc ctgaaatagt ttcaggggaa atgcttggtt
4561 cccgaatcgg atccgcgtat tcgctggatc cctcgatccg ctggtctcta ggtcccgat
4621 gctgcaattc ttacaacagg acttggcata gggtaagcgc aaatgctgtt aaccacacta
4681 acacactttt ttttttcttt tttttttttg agacagagtc tcaactgtgc ggcttggtg
4741 gagtgcagtg gcacgatctc ggctcactgc aacctccggc tccccggctc aagcaattct
4801 cctgcctcag cctcccgagt agctgggatt acaggcatgt gccaccacgc ccggtcaatt
4861 tttgtatttt tagttgagat ggggtttcac catgttggcg aggctggtct tgaactcctg
4921 acctcaggta atccgccagc ctcggcctcc caaagtgtct ggattacaag cgtgagccac
4981 cgtgcccggc caacagtttt taaatctgtg gagacttcat tcccttgat gccttgagc
5041 cgcgcagact acaactccca tcatgcctgg cagccgctgg ggccgcgatt ccgcacgtcc
5101 cttaccgctc tcaactagtc cggcattctt cgctgttttc ctaactcgcc cgcttgacta
5161 gcgccttgga acagccattt gggctcgtgga gtgcgagcac ggccggccaa tgcgcgagtc
5221 agaggggccag gaggggcgcg gccattcgcc gcccgcccc tgctccgtgg ctggttttct
5281 ccgcgggcgc ctcggggcga acctggagat aatgggcagc acctggggga gccctggctg
5341 ggtgcggctc gctctttgcc tgacgggctt agtgcctcgc ctctacgcgc tgcacgtgaa
5401 ggcgggcgcg gcccgggacc gggattaccg cgcgctctgc gacgtgggca ccgccatcag
5461 ctgttcgcgc gtcttctcct ccagggtgtc acgggagtg gaggcgtgg gcctcgagc
5521 agggcgggcca ggatgccaga tgattattct ggagtctggg atcgggtgtc ccggggaacg
5581 gacacggggc tggactgctc gcggggtcgt tgcacagggg ctgagctacc cagcgatact
5641 ggtgttcgaa ataagagtgc gaggcaagg accagacagt gctggggact gggattatct
5701 cggggactcg cacgtgaatt ggatgccaag gaataacggt gaccaggaaa ggcggggagg
5761 caggatggcg gtagagattg acgatggtct caaggacggc gcgcagggtg aggggggtgt
5821 tggcgatggc tgcgcccagg aacaagggtg cccggtctgg ctgtgcgtga tggccaggcg
5881 ttagcataat gacggaatac agaggaggcg agtgagtggc cagggagctg gagattctcg
5941 ggtccagggc aaagataatc tgcccccgac tcccagctc tgatgcaaaa ccgagtgaac
6001 cgttatacca gccttgccat tttagaatt acttaagggc cgggcgcggt ggccactcc
6061 tgtaatccca gcactttggg aggcggaggg ggatggatca ctgaaagtc ggagttgacc
6121 agccttgcca acatggtgaa agcctgtctc taccaaaaat agaaaaatta atcgggcgct
6181 atggcggtg ccttaatccc agctactcgg gggggctaag gcaggagaat cgcttgaacc
6241 cgggaggcgg aggtttcagt gagccgagat cgcgccactg cactccagcc tgggcccagag
6301 tgagactcgc tctcaaaaaa aaaaaaaaaa aaaaaaaaaa agagacttac ttaaggctca
6361 agatgaaaaa cagggcctac ggagtagcca cgtccggggc tggctctggg agaggggagg
6421 atagggtcag tgacatggaa tcctgacgtg gccaaaagtg cccggtgcca ggagatcatc

```

6481 gacccttgga ctaggatggg aggtcgggga acagaggata gccaggtgg cttcttgga
6541 atcaccttcc tcgggcaggg tccaaggcac tgggttgaca gtcctaacct gggtccaccc
6601 caccacaccc ctctgccagg tggggcaggg gtttcgggct ggtggagcat gtgctgggac
6661 aggacagcat cctcaatcaa tccaacagca tattcgggtg catcttctac aactacagc
6721 tattgttagg tgagtggctc cggccctcc ctgcccgccc cggcccgccc ctcatccccc
6781 ttggctcagct cagcccccact ccatgcaatc ttggatgacc acacagctga cagccagcta
6841 gctgctcatc acggagcgtc ctgcccgtgg ggatgtgggg aggttaactaa caggagtctt
6901 ttaattgggt taagtactgt tagaggctga agggccctta aagacatcct aggtcccccag
6961 gttttttgtt tgtgtgtgtt ttgagacagg gtctggctct gttgcccata gtgaggtcta
7021 ggatgccctt agtgtgact ggcgtgatct cagttcatgg caacctctgc ctccctgccc
7081 aagggatcct cccaccttag cctcccaagc agctggaatc acaggcgtgc accactatgc
7141 ccagctaatt ttgttttttg tttttttttg gtagagatgg tgtctcgcca tgttgcccag
7201 gctgggtctca agcaatctgt ctgcctcagc ctcccaaagt gctgggggga ttacaggcgt
7261 gagctaccat gcccaccaa caccacgtt ttgtggaaaa gatgccgaaa ttccttttta
7321 aggagaagct gagcatgagc tatcttttgt ctcatctagt gctcagcagg aaaatttgta
7381 tctagtccca taagaacaga gagaggaaac aaggagatgg aagacgatgg cggccaggc
7441 ctgtctgatg ccatatgccg gagatgagac tatccattac cacccttccc agcaggctcc
7501 cagctccctt ttgagtcacc ctccccagct ccagagaagg catcactgag ggaggcccag
7561 caccacgggc ctggctgaca catgggtcag acttgccgga ttattttaag aaattttatt
7621 gctcagaact ttccctccct gggcaatggc aagagcttca gagaccagtc ccttgagggg
7681 gacctgttga agccttcttt tttttttttt ttaagaaata atcttgctct gttgcccagg
7741 ctggagtgca gtggcacaat catagctcac tgtaacctgg ctcaagcgat cctcctgagt
7801 agctaggact ataggcatgt cactgcaccc agctaatttt tttttttttt
7861 ttttgcgaca tagtctcgct ctgtcaccag gctggagtgc agtggcacga tcttggtcga
7921 ctgcaacctc tgcctcccgg gttcaagcaa ttttcctgcc tcagcctcct gtagtctgg
7981 gactacaggc gcgtgtcacc acgcccagct aatttttgta tttttagtgg agacagggtt
8041 tcaccatggt ggctaggatg gtctcaatct cttgacctgg tgatccatcc gccttgacct
8101 cccaaagtgc taggattaca ggcgtgagtc aacctcaccg ggcatttttt ttttgagacg
8161 aagtcttgcct ctgtctgccc aagctggaat gtgggtggcat gatctcggct cactgcaacc
8221 tccacctcct aggttcaagc gattctccac cttagcctcc ccagcagctg ggattacagg
8281 tgcccacaa cacaccgggc taatttttgt atttttatta gagatgggggt tttgcatgt
8341 tggccagggt gctctgaaac tccaaacctc aggtgatcca ccccattgg cctcccaaaa
8401 tactgggatt acaggcatga gccaccgtgc ccagctgaat ttctaaattt ttgatagaga
8461 tcgggtcttt ctatgttggc caagctgggc ttgaactcct agcctaaagc
agtcctccca
8521 cctcgccctc ccagagtgtt tggaatacgt gcgtaagcca ccacatctgc
cctggagcct
8581 cttgttttag agaccctcc cagcagctcc tggcatctag gtagtgcagt
gacatcatgg
8641 agtggtcggg aggtggccag tgctgaagc ccacaccgga cctcttctg
ccttgagggt
8701 tgctgcgga cacgctgggc ctctgtcctg atgtgtctga gtcctcgggt
gtctctcgct
8761 ggttctgtct acctggcctg gatcctgttc ttcgtgtctt atgatttctg
cattgtttgt
8821 atcacacct atgtatcaa cgtgagcctg atgtggctca gtttccggaa
gggtccaagaa
8881 cccaggggca aggttaagag gcaactgagc ctcaacccaa gccaggctga
cctcatctgc
8941 tttgcttttg catgtgagcc ttgcctaagg gggcatatct gggtccctag
aaggccctag
9001 atgtgggggt tctagattac cccctcctcc tgccataccc gcacatgaca
atggaccaaa
9061 tgtgccacac gctcgtcttt ttttacccc agtgcctctg actctgtccc
catgggctgg
9121 tctccaaagc tctttccatt gccaggggag ggaaggttct gagcaataaa
gtttcttaga

9181 tcaatcagcc aagtctgaac catgtgtctg ccatggactg tgggtgctggg
cctccctcgg
9241 tgttgccctc tctggagctg ggaagggtga gtcagaggga gagtggaggg
cctgctggga
9301 aggggtggtta tgggtagtct catctccagt gtgtggagtc agcaaggcct
ggggcaccat
9361 tggcccccac cccagggaaa caggctggca gctcgtcct gctgcccaca
ggagccaggc
9421 ctccctctcct gggaaggctg agcacacacc tggaagggca ggctgccctt
ctggttctgt
9481 aaatgcttgc tgggaagttc ttccttgagt ttaactttaa cccctccagt
tgccttatcg
9541 accattccaa gccagtattg gtagccttgg agggtcaggg ccaggttgtg
aaggtttttg
9601 ttttgccat tatgccctga ccacttacct acatgccaa gactgtttaa
gaacttgtgt
9661 tggcagggtg cagtggctca cacctgtaac ccctgtactt tgggaggcca
aggcaggagg
9721 atcacttgag gccaggagt ccagaccagc ctgggcaaaa tagtgagacc
cctgtctcta
9781 caaaaaaaaa aaaaaaaaaa aattagccag gcatgggtgt gtatgtacct
atagtcccaa
9841 ctaatcggga agctggcggg aagactgctt gagcccagaa ggttgaggct
gcagtgagcc
9901 atgatcactg cactccagcc tgagcaacag agcaagaccg tctccaaaaa
aaaacaaaaa
9961 acaaaaaaaaa acttgtgtta acgtgttaaa ctcttttaac ctttacagt
atttatgagg
10021 tgggtactat tattatccct atcttgatga tagggacaga gtggctaatt
agtatgcctg
10081 agatcacaca gctactgcag gaggctctca ggatttgaat ccacctggtc
catctggctc
10141 cagcatctat atgctttttt ttttgttggg ttgtttttga gacggacttt
cgctcttggt
10201 gtccaggctg gagtgcattg gcacaatttc ggctcaccac aacctccacc
tcccagggtc
10261 aagtgattct cctgcctcag cctcccaggt agctgggatt acaggcatgc
gccaccaggc
10321 ctggctaatt ttgtattttt agtagagaca gtgtttctct atgttgggtca
ggctgggtctc
10381 gcaactccga cctcagggtga tctgcctgct ttggcctccc aaagtgtctg
gattacaggc
10441 gtgagccacc atgccggcc agcatctata atatatgcta aaccttacc
acagctgcct
10501 ctgtcttggg gtcagggttc ttgaatggtc tgggccagcc tgagttagga
gtgggaagag
10561 gaagaggagg cacgcgcaa agtgttcttc ccgcctgacc gagtctctg
cgctatgag
10621 gaaggtgggc accgtgtca ccctgtttcc cagggaagcca gcaggcccag
ggagatactg
10681 ggatttgcca aaggtcaca gcgagcaagc agcagtgggt gctggctcct
caccactcag
10741 cctggcacac actgtctctg agcagacctg ggagaaggca tgtgaaacct
acctagccct
10801 ggctggaaga gaagtggggg agtggaaatg aaaagtgggg actgggagga
gctgccaccg

10861 tgcctccat ggttcagga tggcactttc atgctgggag tgagcttgcc
ctgtgccagg
10921 ctgtcattga gacgttttcc attcaacaac tgtttctcag acaaggcaca
ggcagggttag
10981 ctctcccagg ccactcaagt agtaaggaaa cagatttaaa cccagacaat
ttggctccac
11041 ggctaaggcc cagagagggg tggtgacaag cctaggggtca cacagccaca
gccccagagc
11101 taggaactgc tggctagggc aagcagacag ctgaatccta gccaaactgga
ggtggagcag
11161 aggtggagcc agggagaccc aggtggcctc

Figure 3

